

CONTROL ID: 1482252

TITLE: Combined Sr, Nd, Pb and Hf isotopic constraints on the origin of Shatsky Rise (NW Pacific)

AUTHORS (FIRST NAME, LAST NAME): Joerg Geldmacher¹, Ken Heydolph¹, David T Murphy², Irina Hoernle¹

INSTITUTIONS (ALL): 1. Dynamics of the Ocean Floor, GEOMAR - Helmholtz Centre for Ocean Research
2. School of Natural Resource Sciences, Queensland University of Technology, Gardens Point, QLD, Australia
3. Geology and Geophysics, School of Ocean and Earth Science and Technology, University of Hawaii, HI, USA

ABSTRACT BODY: The submarine Shatsky Rise plateau in the northwest Pacific Ocean (ca. 1500 km²) is a volcanic plateau of Jurassic to Early Cretaceous age. Based on magnetic reversals combined with bathymetric data, the three main volcanic massifs are proposed to have successively formed by massive volcanism along a southwest-northeast moving spreading center. We investigate a proposed interaction of a possible mantle plume head with the spreading system, Shatsky Rise volcano, in 2009 (Expedition 324 Scientists, 2010). Based on major and trace element compositions, the origin of the lavas can be explained by derivation from a normal mid-ocean ridge basalt (MORB)-like source, although a distinct depletion in Nb and Ta implies that melting started at greater depth (Sano et al. in press). A small fraction of samples (all from Ori massif) show highly over moderately incompatible trace element ratios indicating an enriched (plume?) source.

We present compiled Sr, Nd, Pb and Hf isotope ratios from all three volcanic edifices of Shatsky Rise and compare them to a trace element study. Most isotope data overlap with Pacific MORB composition although regional variations are observed. Drill sites on the oldest edifice, Tamu massif, yield fairly uniform compositions, a wider spread is found for lavas from Ori and Shirshov, suggesting that the source has become more heterogeneous with time (also consistent with the observed trends to reflect a decreasing degree of melting (and therefore less homogenization of inherent plume heterogeneities) during the interaction of the spreading center with a waning plume head. Interestingly, lavas from the Ori massif show elevated ¹⁴³Nd/¹⁴⁴Nd and ²⁰⁶Pb/²⁰⁴Pb ratios, placing them closer to a composition recently proposed for a non-choyudo source which is supposed to be preferentially sampled by large igneous province volcanism (Jackson and Carlson, 2011).

Expedition 324 Scientists, IODP Prel. Rept. 324, 2010

Jackson, M.G. and Carlson, R.W., Nature, vol. 476, 2011

Sano T., Shimizu K., Ishikawa A., Senda R., Chang Q., Kimura J.-I., Widdowson M., Sager W.W. Geochemistry, 2011

KEYWORDS: [1040] GEOCHEMISTRY / Radiogenic isotope geochemistry, [1021] GEOCHEMISTRY / Oceanic crust, [1022] GEOCHEMISTRY / TECTONOPHYSICS / Hotspots, large igneous provinces, and flood basalt volcanism.

(No Image Selected)

(No Table Selected)

Additional Details

Previously Presented Material: Some preliminary data (30%) were presented at last years AGU fall meeting

Contact Details

CONTACT (NAME ONLY): Joerg Geldmacher

CONTACT (E-MAIL ONLY): jgeldmacher@geomar.de

TITLE OF TEAM: