



Seismic structure of lithosphere emplaced at ultra-slow spreading rates

Ingo Grevemeyer (1), Michaela Merz (1), Anke Dannowski (1), Cord Papenberg (1), Nicholas Hayman (2), Harm van Avendonk (2), and Christine Peirce (3)

(1) GEOMAR Helmholtz Centre of Ocean Research, RD4 - Marine Geodynamics, Kiel, Germany (igrevemeyer@geomar.de), (2) Institute for Geophysics, University of Texas at Austin, Texas, U.S.A., (3) Department of Earth Sciences, Durham University, Durham, U.K.

About 57% of the Earth's surface is covered by oceanic crust and new ocean floor is continuously created along the ~60.000 km long mid-ocean ridge (MOR) system. About 25% of the MOR spread at an ultra-slow spreading rate of <20 mm/yr. At ultra-slow spreading rates the melt supply to the ridge is thought to dramatically decrease and crustal thickness decreases to a thickness of <6 km. However, we know little about the processes shaping crust at reduced spreading rates. A formation of crust from a magma chamber would suggest the creation of a well stratified crust, with an extrusive upper crust (layer 2) and a lower gabbroic crust (lower 3) and a well-defined crust-mantle boundary and hence a seismic Moho. In contrast, decompressional melting without formation of a magma chamber would support a crustal structure where seismic velocities change gradually from values typical of crustal rocks to mantle rocks. Here, we report results from a survey from the ultra-slow spreading Cayman Spreading Centre in the Caribbean Sea, sampling mature crust along a flowline from both conjugated ridge flanks. The seismic refraction and wide-angle survey was conducted using ocean-bottom-seismometers from Germany, the UK, and Texas and a 5500 cubic-inch airgun-array source towed by the German research vessel METEOR in April 2015. Typical crustal P-wave velocities support a thin crust of 3 to 5 km thickness. However, a well-defined Moho boundary was not observed. Thus, velocities change gradually from crustal-type velocities (<7.2 km/s) to values of 7.5-7.8 km/s, supporting mantle rocks. In addition to P-waves, we have sampled S-waves along the profile, yielding for the first time V_p/V_s ratios for lithosphere emplaced at ultra-slow spreading rates. Interestingly, about 15 to 20% of the lithosphere has V_p/V_s ratios of >1.9, supporting serpentine. Domains of high V_p/V_s ratio also occur right at the seafloor, supporting large-scale exposure of mantle as proposed by geological evidence from ultra-slow spreading ridges.