



Effect of ocean acidification and elevated $f\text{CO}_2$ on trace gas production by a Baltic Sea summer phytoplankton community

Alison L. Webb^{1,2}, Emma Leedham-Elvidge¹, Claire Hughes³, Frances E. Hopkins⁴, Gill Malin¹, Lennart T. Bach⁵, Kai Schulz⁶, Kate Crawford⁷, Corina P. D. Brussaard^{7,8}, Annegret Stuhr⁵, Ulf Riebesell⁵, and Peter S. Liss¹

¹Centre for Ocean and Atmospheric Sciences, School of Environmental Science, University of East Anglia, Norwich, NR4 7TJ, UK

²Groningen Institute for Evolutionary Life Sciences, University of Groningen, 9700 CC Groningen, the Netherlands

³Environmental Department, University of York, York, YO10 5DD, UK

⁴Plymouth Marine Laboratory, Plymouth, PL1 3DH, UK

⁵GEOMAR Helmholtz Centre for Ocean Research Kiel, Düsternbrooker Weg 20, 24148 Kiel, Germany

⁶Centre for Coastal Biogeochemistry, School of Environment, Science and Engineering, Southern Cross University, Lismore, NSW 2480, Australia

⁷Department of Biological Oceanography, NIOZ – Royal Netherlands Institute for Sea Research, P.O. Box 59, 1790 AB Den Burg, Texel, the Netherlands

⁸Aquatic Microbiology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, P.O. Box 94248, 1090 GE Amsterdam, the Netherlands

Correspondence to: Alison L. Webb (a.l.webb@rug.nl)

Received: 6 November 2015 – Published in Biogeosciences Discuss.: 28 January 2016

Revised: 17 June 2016 – Accepted: 12 July 2016 – Published: 15 August 2016

Abstract. The Baltic Sea is a unique environment as the largest body of brackish water in the world. Acidification of the surface oceans due to absorption of anthropogenic CO_2 emissions is an additional stressor facing the pelagic community of the already challenging Baltic Sea. To investigate its impact on trace gas biogeochemistry, a large-scale mesocosm experiment was performed off Tvärminne Research Station, Finland, in summer 2012. During the second half of the experiment, dimethylsulfide (DMS) concentrations in the highest- $f\text{CO}_2$ mesocosms (1075–1333 μatm) were 34 % lower than at ambient CO_2 (350 μatm). However, the net production (as measured by concentration change) of seven halocarbons analysed was not significantly affected by even the highest CO_2 levels after 5 weeks' exposure. Methyl iodide (CH_3I) and diiodomethane (CH_2I_2) showed 15 and 57 % increases in mean mesocosm concentration (3.8 ± 0.6 increasing to $4.3 \pm 0.4 \text{ pmol L}^{-1}$ and 87.4 ± 14.9 increasing to $134.4 \pm 24.1 \text{ pmol L}^{-1}$ respectively) during Phase II of the experiment, which were unrelated to CO_2 and corresponded to 30 % lower Chl *a* concentrations compared to Phase I. No other iodocarbons increased or showed

a peak, with mean chloriodomethane (CH_2ClI) concentrations measured at $5.3 (\pm 0.9) \text{ pmol L}^{-1}$ and iodoethane ($\text{C}_2\text{H}_5\text{I}$) at $0.5 (\pm 0.1) \text{ pmol L}^{-1}$. Of the concentrations of bromoform (CHBr_3 ; mean $88.1 \pm 13.2 \text{ pmol L}^{-1}$), dibromomethane (CH_2Br_2 ; mean $5.3 \pm 0.8 \text{ pmol L}^{-1}$), and dibromochloromethane (CHBr_2Cl , mean $3.0 \pm 0.5 \text{ pmol L}^{-1}$), only CH_2Br_2 showed a decrease of 17 % between Phases I and II, with CHBr_3 and CHBr_2Cl showing similar mean concentrations in both phases. Outside the mesocosms, an upwelling event was responsible for bringing colder, high- CO_2 , low-pH water to the surface starting on day *t*16 of the experiment; this variable CO_2 system with frequent upwelling events implies that the community of the Baltic Sea is acclimated to regular significant declines in pH caused by up to 800 μatm $f\text{CO}_2$. After this upwelling, DMS concentrations declined, but halocarbon concentrations remained similar or increased compared to measurements prior to the change in conditions. Based on our findings, with future acidification of Baltic Sea waters, biogenic halocarbon emissions are likely to remain at similar values to today; however, emissions of biogenic sulfur could significantly decrease in this region.

1 Introduction

Anthropogenic activity has increased the fugacity of atmospheric carbon dioxide ($f\text{CO}_2$) from 280 μatm (pre-Industrial Revolution) to over 400 μatm today (Hartmann et al., 2013). The IPCC AR5 long-term projections for atmospheric $p\text{CO}_2$ and associated changes to the climate have been established for a variety of scenarios of anthropogenic activity until the year 2300. As the largest global sink for atmospheric CO_2 , the global ocean has absorbed an estimated 30 % of excess CO_2 produced (Canadell et al., 2007). With atmospheric $p\text{CO}_2$ projected to possibly exceed 2000 μatm by the year 2300 (Collins et al., 2013; Cubasch et al., 2013), the ocean will take up increasing amounts of CO_2 , with a potential lowering of surface ocean pH by over 0.8 units (Raven et al., 2005). The overall effect of acidification on the biogeochemistry of surface ocean ecosystems is unknown and currently unquantifiable, with a wide range of potential positive and negative impacts (Doney et al., 2009; Hofmann et al., 2010; Ross et al., 2011).

A number of volatile organic compounds are produced by marine phytoplankton (Liss et al., 2014), including the climatically important trace gas dimethylsulfide (DMS, $\text{C}_2\text{H}_6\text{S}$) and a number of halogen-containing organic compounds (halocarbons), including methyl iodide (CH_3I) and bromoform (CHBr_3). These trace gases are a source of sulfate particles and halide radicals when oxidised in the atmosphere and have important roles as ozone catalysts in the troposphere and stratosphere (O'Dowd et al., 2002; Solomon et al., 1994) and as cloud condensation nuclei (CCNs; Charlson et al., 1987).

DMS is found globally in surface waters originating from the algal-produced precursor dimethylsulfoniopropionate (DMSP; $\text{C}_5\text{H}_{10}\text{O}_2\text{S}$). Both DMS and DMSP provide the basis for major routes of sulfur and carbon flux through the marine microbial food web and can provide up to 100 % of the bacterial and phytoplanktonic sulfur demand (Simó et al., 2009; Vila-Costa et al., 2006a). DMS is also a volatile compound which readily passes through the marine boundary layer to the troposphere, where oxidation results in a number of sulfur-containing particles important for atmospheric climate feedbacks (Charlson et al., 1987; Quinn and Bates, 2011); for this reason, any change in the production of DMS may have significant implications for climate regulation. Several previous acidification experiments have shown differing responses of both compounds (e.g. Avgoustidi et al., 2012; Hopkins et al., 2010; Webb et al., 2015), while others have shown delayed or more rapid responses as a direct effect of CO_2 (e.g. Archer et al., 2013; Vogt et al., 2008). Further, some laboratory incubations of coastal microbial communities showed increased DMS production with increased $f\text{CO}_2$ (Hopkins and Archer, 2014) but lower DMSP production. The combined picture arising from existing studies is that the response of communities to $f\text{CO}_2$ perturbation is not predictable and requires further study. Previous studies mea-

suring DMS in the Baltic Sea measured concentrations up to 100 nmol L^{-1} during the summer bloom, making the Baltic Sea a significant source of DMS (Orlikowska and Schulz-Bull, 2009).

In surface waters, halocarbons such as methyl iodide (CH_3I), chloriodomethane (CH_2ClI), and bromoform (CHBr_3) are produced by biological and photochemical processes: many marine microbes (for example cyanobacteria: Hughes et al., 2011; diatoms: Manley and De La Cuesta, 1997; and haptophytes: Scarratt and Moore, 1998) and macroalgae (e.g. brown-algal *Fucus* species: Chance et al., 2009; red algae: Leedham et al., 2013) utilise halides from seawater and emit a range of organic and inorganic halogenated compounds. This production can lead to significant annual flux to the marine boundary layer in the order of 10 Tg iodine-containing compounds (“iodocarbons”; O'Dowd et al., 2002) and 1 Tg bromine-containing compounds (“bromocarbons”; Goodwin et al., 1997) into the atmosphere. The effect of acidification on halocarbon concentrations has received limited attention, but two acidification experiments measured lower concentrations of several iodocarbons, while bromocarbons were unaffected by $f\text{CO}_2$ up to 3000 μatm (Hopkins et al., 2010; Webb, 2015), whereas an additional mesocosm study did not elicit significant differences from any compound up to 1400 μatm $f\text{CO}_2$ (Hopkins et al., 2013).

Measurements of the trace gases within the Baltic Sea are limited, with no prior study of DMSP concentrations in the region. The Baltic Sea is the largest body of brackish water in the world, and salinity ranges from 1 to 15. Furthermore, seasonal temperature variations of over 20 °C are common. A permanent halocline at 50–80 m separates CO_2 -rich, bottom waters from fresher, lower- CO_2 surface waters, and a summer thermocline at 20 m separates warmer surface waters from those below 4 °C (Janssen et al., 1999). Upwelling of bottom waters from below the summer thermocline is a common summer occurrence, replenishing the surface nutrients while simultaneously lowering surface temperature and pH (Brutemark et al., 2011). Baltic organisms are required to adapt to significant variations in environmental conditions. The species assemblage in the Baltic Sea is different to those studied during previous mesocosm experiments in the Arctic, North Sea, and Korea (Brussaard et al., 2013; Engel et al., 2008; Kim et al., 2010) and is largely unstudied in terms of its community trace gas production during the summer bloom. Following the spring bloom (July–August), a low dissolved inorganic nitrogen (DIN) to dissolved inorganic phosphorous (DIP) ratio combines with high temperatures and light intensities to encourage the growth of heterocystous cyanobacteria (Niemisto et al., 1989; Raateoja et al., 2011), in preference to nitrate-dependent groups.

Here we report the concentrations of DMS, DMSP, and halocarbons from the 2012 summer post-bloom season mesocosm experiment aimed to assess the impact of elevated $f\text{CO}_2$ on the microbial community and trace gas pro-

duction in the Baltic Sea. Our objective was to assess how changes in the microbial community driven by changes in $f\text{CO}_2$ impacted DMS and halocarbon concentrations. It is anticipated that any effect of CO_2 on the growth of different groups within the phytoplankton assemblage will result in an associated change in trace gas concentrations measured in the mesocosms as $f\text{CO}_2$ increases, which can potentially be used to predict future halocarbon and sulfur emissions from the Baltic Sea region.

2 Methods

2.1 Mesocosm design and deployment

Nine mesocosms were deployed on the 10 June 2012 (day $t-10$; days are numbered negative prior to CO_2 addition and positive afterward) and moored near Tvärminne Zoological Station ($59^\circ 51.5' \text{ N}$, $23^\circ 15.5' \text{ E}$) in Tvärminne Storfjärden in the Baltic Sea. Each mesocosm comprised a thermoplastic polyurethane (TPU) enclosure of 17 m depth, containing approximately 54 000 L of seawater, supported by an 8 m tall floating frame capped with a polyvinyl hood. For full technical details of the mesocosms, see Czerny et al. (2013) and Riebesell et al. (2013). The mesocosm bags were filled by lowering through the stratified water column until fully submerged, with the opening at both ends covered by 3 mm mesh to exclude organisms larger than 3 mm such as fish and large zooplankton. The mesocosms were then left for 3 days ($t-10$ to $t-7$) with the mesh in position to allow exchange with the external water masses and ensure the mesocosm contents were representative of the phytoplankton community in the Storfjärden. On $t-7$, the bottom of the mesocosm was sealed with a sediment trap and the upper opening was raised to approximately 1.5 m above the water surface. Stratification within the mesocosm bags was broken up on $t-5$ by the use of compressed air for 3.5 min to homogenise the water column and ensure an even distribution of inorganic nutrients at all depths. Unlike in previous experiments, there was no addition of inorganic nutrients to the mesocosms at any time during the experiment; mean inorganic nitrate, inorganic phosphate, and ammonium concentrations measured across all mesocosms at the start of the experiment were $37.2 (\pm 18.8 \text{ SD})$, $323.9 (\pm 19.4 \text{ SD})$, and $413.8 (\pm 319.5 \text{ SD}) \text{ nmol L}^{-1}$ respectively.

To obtain mesocosms with different $f\text{CO}_2$, the carbonate chemistry of the mesocosms was altered by the addition of different volumes of 50 μm filtered, CO_2 -enriched Baltic Sea water (sourced from outside the mesocosms), to each mesocosm over a 4-day period, with the first day of addition being defined as day $t0$. The addition of the enriched CO_2 water was by the use of a bespoke dispersal apparatus (“Spider”) lowered through the bags to ensure even distribution throughout the water column (further details are in Riebesell et al., 2013). Measurements of salinity in the meso-

cosms throughout the experiment determined that three of the mesocosms were not fully sealed and had undergone unquantifiable water exchange with the surrounding waters. These three mesocosms (M2, M4, and M9) were excluded from the analysis. Two mesocosms were designated as controls (M1 and M5) and received only filtered seawater via the Spider; four mesocosms received the addition of CO_2 -enriched waters, with the range of target $f\text{CO}_2$ levels between 600 and 1650 μatm (M7, 600; M6, 950; M3, 1300; M8 1650 μatm). Mesocosms were randomly allocated a target $f\text{CO}_2$; a noticeable decrease in $f\text{CO}_2$ was identified in the three highest- $f\text{CO}_2$ mesocosms (M6, M3, and M8) over the first half of the experiment, which required the addition of more CO_2 -enriched water on $t15$ to bring the $f\text{CO}_2$ back up to maximum concentrations (Fig. 1a; Paul et al., 2015). A summary of the $f\text{CO}_2$ in the mesocosms can be seen in Table 1. At the same time as this further CO_2 addition on $t15$, the walls of the mesocosms were cleaned using a bespoke wiper apparatus (see Riebesell et al., 2013, for more information), followed by weekly cleaning to remove aggregations on the film which would block incoming light. Light measurements showed that over 95 % of the photosynthetically active radiation (PAR) was transmitted by the clean TPU and PVC materials with 100 % absorbance of UV light (Riebesell et al., 2013). Samples for most parameters were collected from the mesocosms at the same time every morning from $t-3$ and analysed daily or every other day.

2.2 Trace gas extraction and analysis

2.2.1 DMS and halocarbons

A depth-integrated water sampler (IWS, HYDRO-BIOS, Kiel, Germany) was used to sample the entire 17 m water column daily or every other day. As analysis of chlorophyll *a* (Chl *a*) showed it to be predominantly produced in the first 10 m of the water column, trace gas analysis was conducted only on integrated samples collected from the surface 10 m, with all corresponding community parameter analyses with the exception of pigment analysis performed also to this depth. Water samples for trace gas analysis were taken from the first IWS from each mesocosm to minimise the disturbance and bubble entrainment from taking multiple samples in the surface waters. As in Hughes et al. (2009), samples were collected in 250 mL amber glass bottles in a laminar flow with minimal disturbance to the water sample, using Tygon tubing from the outlet of the IWS. Bottles were rinsed twice before being carefully filled from the bottom with minimal stirring and allowed to overflow the volume of the bottle approximately three times before sealing with a glass stopper to prevent bubble formation and atmospheric contact. Samples were stored below 10°C in the dark for 2 h prior to analysis. Each day, a single sample was taken from each mesocosm, with two additional samples taken from one randomly

Table 1. Summary of $f\text{CO}_2$ and pH_T (total scale) during phases 0, I, and II of the mesocosm experiment.

Mesocosm*	Target $f\text{CO}_2$ (μatm)	Whole experiment ($t-3$ to $t31$)		Phase 0 ($t-3$ to $t0$)		Phase I ($t1$ to $t16$)		Phase II ($t16$ to $t31$)	
		Mean $f\text{CO}_2$ (μatm)	Mean pH_T						
M1	Control	331	7.91	231	8.00	328	7.95	399	7.86
M5	Control	334	7.91	244	7.98	329	7.94	399	7.52
M7	390	458	7.80	239	7.99	494	7.81	532	7.76
M6	840	773	7.63	236	7.99	932	7.59	855	7.59
M3	1120	950	7.56	243	7.98	1176	7.51	1027	7.52
M8	1400	1166	7.49	232	8.00	1481	7.43	1243	7.45
Baltic Sea	380	350	7.91	298	7.91	277	7.98	436	7.86

* Listed in order of increasing $f\text{CO}_2$.

selected mesocosm to evaluate the precision of the analysis ($<4\%$, no further data shown).

On return to the laboratory, 40 mL of water was injected into a purge and cryotrap system (Chuck et al., 2005), filtered through a 25 mm Whatman glass fibre filter (GF/F; GE Healthcare Life Sciences, Little Chalfont, England) and purged with oxygen-free nitrogen (OFN) at 80 mL min^{-1} for 10 min. Each gas sample passed through a glass wool trap to remove particles and aerosols, before a dual Nafion counterflow drier (180 mL min^{-1} OFN) removed water vapour from the gas stream. The gas sample was trapped in a stainless steel loop held at -150°C in the headspace of a liquid-nitrogen-filled dewar. The sample was injected by immersion of the sample loop in boiling water into an Agilent 6890 gas chromatograph equipped with a 60 m DB-VRX capillary column (0.32 mm ID, $1.8\ \mu\text{m}$ film thickness, Agilent J&W Ltd) according to the programme outlined by Hopkins et al. (2010). Analysis was performed by an Agilent 5973 quadrupole mass spectrometer operated in electron ionisation, single-ion mode. Liquid standards of CH_3I , diiodomethane (CH_2I_2), CH_2ClI , iodoethane ($\text{C}_2\text{H}_5\text{I}$), iodopropane ($\text{C}_3\text{H}_7\text{I}$), CHBr_3 , dibromoethane (CH_2Br_2), dibromochloromethane (CHBr_2Cl), bromiodomethane (CH_2BrI), and DMS (standards supplied by Sigma Aldrich Ltd, UK) were gravimetrically prepared by dilution in high-performance liquid chromatography (HPLC) grade methanol (Table 2) and used for calibration. The relative standard error was expressed as a percentage of the mean for the sample analysis, calculated for each compound using triplicate analysis each day from a single mesocosm, and was $<7\%$ for all compounds. Gas chromatography–mass spectrometry instrument drift was corrected by the use of a surrogate analyte standard in every sample, comprising deuterated DMS ($\text{D}_6\text{-DMS}$), deuterated methyl iodide (CD_3I), and ^{13}C dibromoethane ($^{13}\text{C}_2\text{H}_4\text{Br}_2$) via the method described in Hughes et al. (2006) and Martino et al. (2005). Five-point calibrations were performed weekly for each compound with

Table 2. Calibration ranges and calculated percentage mean relative standard error for the trace gases measured in the mesocosms.

Compound	Calibration range (pmol L^{-1})	% Mean relative standard error
DMS	600–29 300*	6.33
DMSP	2030–405 900*	
CH_3I	0.11–11.2	4.62
CH_2I_2	5.61–561.0	4.98
$\text{C}_2\text{H}_5\text{I}$	0.10–4.91	5.61
CH_2ClI	1.98–99.0	3.64
CHBr_3	8.61–816.0	4.03
CH_2Br_2	0.21–20.9	5.30
CHBr_2Cl	0.07–7.00	7.20

* Throughout the rest of this paper, these measurements are given in nmol L^{-1} .

the addition of the surrogate analyte, with a single standard analysed daily to check for instrument drift; linear regression from calibrations typically produced $r^2 > 0.98$. All samples measured within the mesocosms were within the concentration ranges of the calibrations (Table 2).

2.2.2 DMSP

Samples for total DMSP (DMSP_T) were collected and stored for later analysis by the acidification method of Curran et al. (1998). A 7 mL subsample was collected from the amber glass bottle into an 8 mL glass sample vial (Labhut, Churham, UK), into which $0.35\ \mu\text{L}$ of 50 % H_2SO_4 was added, before storage at ambient temperature. Particulate DMSP (DMSP_P) samples were prepared by the gravity filtration of 20 mL of sample through a 47 mm GF/F in a glass filter unit, before careful removal and folding of the GF/F into a 7 mL sample vial filled with 7 mL of Milli-Q water and $0.35\ \mu\text{L}$ of H_2SO_4 before storage at ambient temperature. Samples were stored for approximately 8 weeks prior




Figure 1. Daily measurements of (a) $f\text{CO}_2$, (b) mean temperature, and (c) total chlorophyll a in the mesocosms and surrounding Baltic Sea waters. Dashed lines represent the three phases of the experiment, based on the Chl a data.

to analysis. DMSP samples (total and particulate) were analysed on a PTFE purge and cryotrap system using 2 mL of the sample purged with 1 mL of 10 M NaOH for 5 min at 80 mL min^{-1} . The sample gas stream passed through a glass wool trap and Nafion counterflow (Permapure) drier before being trapped in a PTFE sample loop kept at -150°C by suspension in the headspace of a liquid nitrogen-filled dewar and controlled by feedback from a thermocouple. Immersion in boiling water rapidly re-volatilised the sample for injection into a Shimadzu GC2010 gas chromatograph with a Varian Chrompack CP-Sil-5CB column (30 m, 0.53 mm ID)

and flame photometric detector (FPD). The gas chromatography (GC) oven was operated isothermally at 60°C , which resulted in DMS eluting at 2.1 min. Liquid DMSP standards were prepared and purged in the same manner as the sample to provide weekly calibrations of the entire analytical system. Involvement in the 2013 AQA 12-23 international DMS analysis proficiency test (National Measurement Institute of Australia, 2013) in February 2013 demonstrated excellent agreement between our method of DMSP analysis and the mean from 13 laboratories measuring DMS using different methods, with a measurement error of 5 %.

DMSP was not detected in any of the samples (total or particulate) collected and stored during the experiment, and it was considered likely that this was due to an unresolved issue regarding acidifying Baltic Sea samples for later DMSP analysis. This method had been used during a previous mesocosm experiment (SOPRAN II, Bergen, Norway), and the results correlated well with those measured immediately on a similar GC-FPD system (Webb et al., 2015). It was considered unlikely that rates of bacterial DMSP turnover through demethylation rather than through cleavage to produce DMS (Curson et al., 2011) were sufficiently high in the Baltic Sea to remove all detectable DMSP yet still produce measurable DMS concentrations. Also, rapid turnover of dissolved DMSP in surface waters being the cause of low DMSP_T concentrations does not explain the lack of intracellular particulate-phase DMSP. Although production of DMS is possible from alternate sources, it is highly unlikely that there was a total absence of DMSP-producing phytoplankton within the mesocosms or Baltic Sea surface waters around Tvärminne; DMSP was measured in surface waters of the southern Baltic Sea at 22.2 nmol L^{-1} in 2012, indicating that DMSP-producing species are present within the Baltic Sea (C. Zindler, personal communication, 2014).

A previous study by del Valle et al. (2011) highlighted up to 94 % loss of DMSP_T from acidified samples of colonial *Phaeocystis globosa* culture and field samples dominated by colonial *Phaeocystis antarctica*. Despite filamentous, colonial cyanobacteria in the samples from Tvärminne mesocosms potentially undergoing the same process, these species did not dominate the community, at only 6.6 % of the total Chl a , implying that the acidification method for DMSP fixation also failed for unicellular phytoplankton species. The findings of this mesocosm study suggest that the acidification method is unreliable in the Baltic Sea and should be considered inadequate as the sole method of DMSP fixation in future experiments in the region. The DMSP acidification method is used worldwide as a simple and effective method of DMSP storage; the findings here, alongside those of del Valle et al. (2011), question the applicability of this method in other marine environments and suggest significant testing prior to reliance on this method as a sole means of DMSP storage.

2.3 Measurement of carbonate chemistry and community dynamics

Water samples were collected from the 10 and 17 m IWS on a daily basis and analysed for carbonate chemistry, fluorometric Chl *a*, phytoplankton pigments (17 m IWS only), and cell abundance to analyse the community structure and dynamics during the experiment. The carbonate system was analysed through a suite of measurements (Paul et al., 2015), including potentiometric titration for total alkalinity (TA), infrared absorption for dissolved inorganic carbon (DIC), and spectrophotometric determination for pH. For Chl *a* analysis and pigment determination, 500 mL subsamples were filtered through a GF/F and stored frozen (-20°C for 2 h for Chl *a* and -80°C for up to 6 months for pigments), before homogenisation in 90 % acetone with glass beads. After centrifuging (10 min at 800 g at 4°C) the Chl *a* concentrations were determined using a Turner AU-10 fluorometer by the methods of Welschmeyer (1994), and the phytoplankton pigment concentrations were determined by reverse phase high-performance liquid chromatography (WATERS HPLC with a Varian Microsorb-MV 100-3 C8 column) as described by Barlow et al. (1997). Phytoplankton community composition was determined by the use of the CHEMTAX algorithm to convert the concentrations of marker pigments to Chl *a* equivalents (Mackey et al., 1996; Schulz et al., 2013). Microbes were enumerated using a Becton Dickinson FAC-SCalibur flow cytometer (FCM) equipped with a 488 nm argon laser (Crawford et al., 2016), and counts of phytoplankton cells $> 20\ \mu\text{m}$ were made on concentrated (50 mL) sample water, fixed with acidic Lugol's iodine solution with an inverted microscope. Filamentous cyanobacteria were counted in $50\ \mu\text{m}$ length units.

2.4 Statistical analysis

All statistical analysis was performed using Minitab V16. In analysis of the measurements between mesocosms, one-way ANOVA was used with Tukey's post hoc analysis test to determine the effect of different $f\text{CO}_2$ on concentrations measured in the mesocosms and the Baltic Sea (H_0 assumes no significant difference in the mean concentrations of trace gases measured through the duration of the experiment). Spearman's rank correlation coefficients were calculated to compare the relationships between trace gas concentrations, $f\text{CO}_2$, and a number of biological parameters, and the resulting ρ values for each correlation are given in Supplement Table S1 for the mesocosms and Table S2 for the Baltic Sea data.




Figure 2. (a) Mean DMS concentrations measured daily in the mesocosms and Baltic Sea from an integrated water sample of the surface 10 m. Dashed lines show the phases of the experiment as given in Fig. 1; $f\text{CO}_2$ shown in the legend is mean $f\text{CO}_2$ across the duration of the experiment. (b) Mean DMS concentrations from each mesocosm during Phase I (crosses) and Phase II (diamonds), for ambient (blue), medium (grey), and high $f\text{CO}_2$ (red), with error bars showing the range of both the DMS and $f\text{CO}_2$.

3 Results and discussion

3.1 Biogeochemical changes within the mesocosms

The mesocosm experiment was split into three phases based on the temporal variation in Chl *a* (Fig. 2; Paul et al., 2015) evaluated after the experiment was completed:

- Phase 0 (days $t-5$ to t_0) – pre- CO_2 addition;
- Phase I (days t_1 to t_{16}) – “productive phase”;
- Phase II (days t_{17} to t_{30}) – temperature-induced autotrophic decline.

3.1.1 Physical parameters

$f\text{CO}_2$ decreased over Phase I in the three highest- $f\text{CO}_2$ mesocosms, mainly through air–sea gas exchange and carbon fixation by phytoplankton (Fig. 1a). All mesocosms still showed distinct differences in $f\text{CO}_2$ levels throughout the experiment (Table 1), and there was no overlap of mesocosm $f\text{CO}_2$ values on any given day, save for the two controls (M1 and M5). The control mesocosm $f\text{CO}_2$ increased through Phase I of the experiment, likely as a result of undersaturation of the water column encouraging dissolution of atmospheric CO_2 (Paul et al., 2015). Salinity in the mesocosms remained constant throughout the experiment at 5.70 ± 0.004 and showed no variation with depth (data not shown but available in Paul et al., 2015). It remained similar to salinity in the Baltic Sea surrounding the mesocosms, which was 5.74 ± 0.14 . Water temperature varied from a low of $8.6 \pm 0.4^\circ\text{C}$ during Phase 0 to a high of $15.9 \pm 2.2^\circ\text{C}$ measured on day $t16$, before decreasing once again (Fig. 1b).

Summertime upwelling events are common and well described (Gidhagen, 1987; Lehmann and Myrberg, 2008) and induce a significant temperature decrease in surface waters; such an event appears to have commenced around $t16$, as indicated by significantly decreasing temperatures inside and out of the mesocosms (Fig. 1b) and increased salinity in the Baltic Sea from 5.5 to 6.1 over the following 15 days to the end of the experiment. Due to the enclosed nature of the mesocosms, the upwelling affected only the temperature and not pH, $f\text{CO}_2$, or the microbial community. However, the temperature decrease after $t16$ was likely to have had a significant effect on phytoplankton growth (and biogenic gas production), explaining the lower Chl a in Phase II.

3.1.2 Community dynamics

Mixing of the mesocosms and redistribution of the nutrients throughout the water column after closure (prior to $t-3$) did not trigger a notable increase in total Chl a in Phase 0 as was identified in previous mesocosm experiments. During Phase I, light availability, combined with increasing water temperatures, favoured the growth of phytoplankton in all mesocosms (Paul et al., 2015) and was unlikely to be a direct result of the CO_2 enrichment, as no difference was identified between enriched mesocosms and controls. Mean Chl a during Phase I was $1.98 (\pm 0.29) \mu\text{g L}^{-1}$ from all mesocosms, decreasing to $1.44 (\pm 0.46) \mu\text{g L}^{-1}$ in Phase II; this decrease was attributed to a temperature-induced decrease in phytoplankton growth rates and higher grazing rates as a result of higher zooplankton reproduction rates during Phase I (Lischka et al., 2015; Paul et al., 2015). Mesocosm Chl a decreased until the end of the experiment on $t31$.

The largest contributors to Chl a in the mesocosms during the summer of 2012 were the chlorophytes and cryptophytes, with up to 40 and 21 % contributions to the Chl a respectively (Table 3; Paul et al., 2015). Significant long-term differences

in abundance between mesocosms developed as a result of elevated $f\text{CO}_2$ in only two groups: picoeukaryotes I showed higher abundance at high $f\text{CO}_2$ ($F = 8.2$, $p < 0.01$; Crawford et al., 2016, and Supplement Fig. S2), as seen in previous mesocosm experiments (Brussaard et al., 2013; Newbold et al., 2012), and picoeukaryotes III showed the opposite trend ($F = 19.6$, $p < 0.01$; Crawford et al., 2016). Temporal variation in phytoplankton abundance was similar between all mesocosms (Figs. S1 and S2).

Diazotrophic, filamentous cyanobacterial blooms in the Baltic Sea are an annual event in summer (Finni et al., 2001), and single-celled cyanobacteria have been found to comprise as much as 80 % of the cyanobacterial biomass and 50 % of the total primary production during the summer in the Baltic Sea (Stal et al., 2003). However, CHEMTAX analysis identified cyanobacteria as contributing less than 10 % of the total Chl a in the mesocosms (Crawford et al., 2016; Paul et al., 2015). These observations were backed up by satellite observations showing reduced cyanobacterial abundance throughout the Baltic Sea in 2012 compared to previous and later years (Oberger, 2013). It was proposed that light availability and surface water temperatures during the summer of 2012 were suboptimal for triggering a filamentous cyanobacteria bloom (Wasmund, 1997).

3.2 DMS and DMSP

3.2.1 Mesocosm DMS

A significant 34 % reduction in DMS concentrations was detected in the high- $f\text{CO}_2$ treatments during Phase II compared to the ambient- $f\text{CO}_2$ mesocosms ($F = 31.7$, $p < 0.01$). Mean DMS concentrations of $5.0 (\pm 0.8)$; range $3.5\text{--}6.8$ nmol L^{-1} in the ambient treatments were compared to $3.3 (\pm 0.3)$; range $2.9\text{--}3.9$ nmol L^{-1} in the 1333 and 1075 μatm mesocosms (Fig. 2a). The primary differences identified were apparent from the start of Phase II on $t17$, after which maximum concentrations were observed in the ambient mesocosms on $t21$. The relationship between DMS and increasing $f\text{CO}_2$ during Phase II was found to be linear (Fig. 2b), a finding also identified in previous mesocosm experiments (Archer et al., 2013; Webb et al., 2015). Furthermore, increases in DMS concentrations under high $f\text{CO}_2$ were delayed by 3 days relative to the ambient- and medium- $f\text{CO}_2$ treatments, a situation which has been observed in a previous mesocosm experiment. This was attributed to small-scale shifts in community composition and succession which could not be identified with only a once-daily measurement regime (Vogt et al., 2008). DMS measured in all mesocosms fell within the range 2.7 to 6.8 nmol L^{-1} across the course of the experiment. During Phase I, no difference was identified in DMS concentrations between $f\text{CO}_2$ treatments, with the mean of all mesocosms being $3.1 (\pm 0.2) \text{nmol L}^{-1}$. Concentrations in all mesocosms gradually declined from $t21$ until the end of DMS measurements on $t31$. DMS concentrations

Table 3. Abundance and contributions of different phytoplankton groups to the total phytoplankton community assemblage, showing the range of measurements from total Chl a (Paul et al., 2015), CHEMTAX analysis of derived Chl a (Paul et al., 2015), and phytoplankton abundance (Crawford et al., 2016). Data are split into the range of all the mesocosm measurements and those from the Baltic Sea.

	Mesocosm			Baltic Sea		
	Range Integrated 10 m	Range Integrated 17 m	% contribution to Chl a	Range Integrated 10 m	Range Integrated 17 m	% contribution to Chl a
Chl a	0.9–2.9	0.9–2.6	100	1.3–6.5	1.12–5.5	100
Phytoplankton taxonomy (equivalent chlorophyll $\mu\text{g L}^{-1}$)						
Cyanobacteria		0.01–0.4	8		0.0–0.1	1
Prasinophytes		0.04–0.3	7		0.01–0.3	4
Euglenophytes		0.0–1.6	15		0.0–2.6	21
Dinoflagellates		0.0–0.3	3		0.04–0.6	9
Diatoms		0.1–0.3	7		0.04–0.9	9
Chlorophytes		0.3–2.0	40		0.28–3.1	41
Cryptophytes		0.1–1.4	21		0.1–1.0	15
Small phytoplankton (< 10 μm) abundance (cells mL^{-1})						
Cyanobacteria	55 000–380 000	65 000–470 000		30 000–180 000	30 000–250 000	
Picoeukaryotes I	15 000–10 0000	17 000–111 000		5000–70 000	6100–78 000	
Picoeukaryotes II	700–4000	600–4000		400–3000	460–3700	
Picoeukaryotes III	1000–9000	1100–8500		1000–6000	950–7500	
Nanoeukaryotes I	400–1400	270–1500		200–4000	210–4100	
Nanoeukaryotes II	0–400	4–400		100–1100	60–1300	

measured in the mesocosms and Baltic Sea were comparable to those measured in temperate coastal conditions in the North Sea (Turner et al., 1988), the Mauritanian upwelling (Franklin et al., 2009; Zindler et al., 2012), and the South Pacific (Lee et al., 2010).

The majority of DMS production is presumed to be from DMSP. However, an alternative production route for DMS is available through the methylation of methanethiol (Drotar et al., 1987; Kiene and Hines, 1995; Stets et al., 2004), predominantly identified in anaerobic environments such as freshwater lake sediments (Lomans et al., 1997), salt marsh sediments (Kiene and Visscher, 1987), and microbial mats (Visscher et al., 2003; Zinder et al., 1977). Recent studies have also identified this pathway of DMS production from *Pseudomonas deceptionensis* in an aerobic environment (Carrión et al., 2015), where *P. deceptionensis* was unable to synthesise or catabolise DMSP but was able to enzymatically mediate DMS production from methanethiol (MeSH). The same enzyme has also been identified in a wide range of other bacterial taxa, including the cyanobacterial *Pseudanabaena*, which was identified in the Baltic Sea during this and previous investigations (A. Stühr, personal communication, 2015; Kangro et al., 2007; Nausch et al., 2009). Correlations between DMS and the cyanobacterial equivalent Chl a ($\rho = 0.42$, $p < 0.01$; Fig. S1g) and DMS and single-celled cyanobacteria ($\rho = 0.58$, $p < 0.01$; Fig. S2a) suggest that the methylation pathway may be a potential source of DMS within the Baltic Sea community. In addition to the

methylation pathway, DMS production has been identified from S-methylmethionine (Bentley and Chasteen, 2004), as well as from the reduction of dimethylsulfoxide (DMSO), in both surface and deep waters by bacterial metabolism (Hattton et al., 2004). As these compounds were not measured in the mesocosms, it is impossible to determine whether they were significant sources of DMS.

3.2.2 DMS and community interactions

Throughout Phase I, DMS showed no correlation with any measured variables of biological activity or cell abundance and was unaffected by elevated $f\text{CO}_2$, indicating that measured DMS concentrations were not directly related to the perturbation of the system and associated cellular stress (Sunda et al., 2002). Of the studied phytoplankton groupings, neither the cryptophytes nor chlorophytes as the largest contributors of Chl a were identified as significant producers of DMSP. During Phase II, DMS was negatively correlated with Chl a in the ambient- and medium- $f\text{CO}_2$ mesocosms ($\rho = -0.60$, $p < 0.01$). During Phase II, a significant correlation was seen between DMS and single-celled cyanobacteria identified predominantly as *Synechococcus* ($\rho = 0.53$, $p < 0.01$; Crawford et al., 2016, and Table S1) and picoeukaryotes III ($\rho = 0.75$, $p < 0.01$). The peak in DMS concentrations on $t21$ is unlikely to be a delayed response to the increased Chl a on $t16$ due to the time lag of 7 days. These higher DMS concentrations were likely connected to a peak

Table 4. Concentration ranges of trace gases measured in the mesocosms compared to other open-water ocean acidification experiments, showing the range of concentrations for each gas and the percentage change between the control and the highest- $f\text{CO}_2$ treatment. SOPRAN: Surface Ocean Processes in the Anthropocene; NERC: Natural Environment Research Council; EPOCA: European Project on Ocean Acidification; UKOA: UK Ocean Acidification Research Programme.

	Range $f\text{CO}_2$		DMS	CH ₃ I	CH ₂ I ₂	CH ₂ ClI	CHBr ₃	CH ₂ Br ₂	CH ₂ Br ₂ Cl
	(μatm)		(nmol L^{-1})	(pmol L ⁻¹)					
SOPRAN Tvärminne mesocosm (this study)	346–1333	Range % change	2.7–6.8 –34	2.9–6.4 –0.3	57–202 1.3	3.8–8.0 –11	69–148 –9	4.0–7.7 –3	1.7–3.1 –4
SOPRAN Bergen 2011 Webb et al. (2015)	280–3000	Range % change	0.1–4.9 –60	4.9–32 –37	5.8–321 –48	9.0–123 –27	64–306 –2	6.3–30.8 –4	3.9–14 –6
NERC microbial metagenomics experiment, Bergen (2006), Hopkins et al. (2010)	300–750	Range % change	ND–50 –57	2.0–25 –41	ND–750 –33	ND–700 –28	5.0–80 13	ND–5.5 8	0.2–1.2 22
EPOCA Svalbard 2010 Archer et al. (2013), Hopkins et al. (2013)	180–1420	Range % change	ND–14 –60	0.04–10 NS	0.01–2.5	0.3–1.6 NS	35–151 NS	6.3–33.3 NS	1.6–4.7 NS
UKOA European shelf 2011 Hopkins and Archer (2014)	340–1000	Range % change	0.5–12 225						
Korean mesocosm experiment 2012 Park et al. (2014)	160–830	Range % change	1.0–100 –82						

ND – not detected.
NC – no change.

in dissolved organic carbon (DOC) on $t15$, as well as increasing bacterial abundance during Phase II (Hornick et al., 2016). It is also likely that DMS concentrations increased as a response to the mesocosm wall cleaning which took place on $t16$. The variation in inorganic nutrient concentrations between mesocosms at the start of the experiment did not have an effect on DMS concentrations during Phase I, and by the start of Phase II the variation between mesocosms had decreased.

In previous mesocosm experiments (Archer et al., 2013; Hopkins et al., 2010; Webb et al., 2015), DMS has shown poor correlations with many of the indicators of primary production and phytoplankton abundance, as well as showing the same trend of decreased concentrations in high- $f\text{CO}_2$ mesocosms compared to ambient ones. DMS production is often uncoupled from measurements of primary production in open waters (Lana et al., 2012) and also often from the production of its precursor DMSP (Archer et al., 2009). DMS and DMSP are important sources of sulfur and carbon in the microbial food web for both bacteria and algae (Simó et al., 2002, 2009), and since microbial turnover of DMSP and DMS play a significant role in net DMS production, it is unsurprising that DMS concentrations have shown poor correlation with DMSP-producing phytoplankton groups in past experiments and open waters.

DMS concentrations have been reported to be lower under conditions of elevated $f\text{CO}_2$ compared to ambient controls, in both mesocosm experiments (Table 4) and phytoplankton monocultures (Arnold et al., 2013; Avgoustidi et al., 2012).

However, the varying response of the community within each experiment limits our ability to generalise the response of algal production of DMS and DMSP in all situations due to the characteristic community dynamics of each experiment in specific geographical areas and temporal periods. Previous experiments in the temperate Raunefjorden of Bergen, Norway, showed lower abundance of DMSP-producing algal species, and subsequently of DMSP-dependent DMS concentrations (Avgoustidi et al., 2012; Hopkins et al., 2010; Vogt et al., 2008; Webb et al., 2015). In contrast mesocosm experiments in the Arctic and Korea have shown increased abundance of DMSP producers (Archer et al., 2013; Kim et al., 2010) but lower DMS concentrations, while incubation experiments by Hopkins and Archer (2014) showed lower DMSP production but higher DMS concentrations at high $f\text{CO}_2$. However, in all previous experiments with DMSP as the primary precursor of DMS, elevated $f\text{CO}_2$ had a less marked effect on measured DMSP concentrations than on measured DMS concentrations. Hopkins et al. (2010) suggested that “the perturbation of the system has a greater effect on the processes that control the conversion of DMSP to DMS rather than the initial production of DMSP itself”.

Previous mesocosm experiments have suggested significant links between increased bacterial production through greater availability of organic substrates at high $f\text{CO}_2$ (Engel et al., 2013; Piontek et al., 2013). Further, Endres et al. (2014) identified significant enhanced enzymatic hydrolysis of organic matter with increasing $f\text{CO}_2$, with higher bacterial abundance. Higher bacterial abundance will likely




Figure 3. Mean concentrations (pmol L^{-1}) of (a) CH_3I , (b) $\text{C}_2\text{H}_5\text{I}$, (c) CH_2I_2 , and (d) CH_2ClI taken from a water sample integrated from the surface 10 m. Dashed lines indicate the phases of the experiment, as given in Fig. 2. $f\text{CO}_2$ shown in the legend is mean $f\text{CO}_2$ across the duration of the experiment.

result in greater bacterial demand for sulfur and therefore greater consumption of DMS and conversion to DMSO. This was suggested as a significant sink for DMS in a previous experiment (Webb et al., 2015), but during the present experiment, both bacterial abundance and bacterial production were lower at high $f\text{CO}_2$ (Hornick et al., 2016). However, as it has been proposed that only specialist bacterial groups are DMS consumers (Vila-Costa et al., 2006b) and there is no determination of the DMS consumption characteristics of the bacterial community in the Baltic Sea, it is not known if this loss pathway is stimulated at high $f\text{CO}_2$. As microbial DMS yields can vary between 5 and 40 % depending on the sulfur and carbon demand (Kiene and Linn, 2000), a change in the bacterial sulfur requirements could change DMS turnover despite lower abundance.

3.3 Iodocarbons in the mesocosms and relationships with community composition

Elevated $f\text{CO}_2$ did not affect the concentration of iodocarbons in the mesocosms significantly at any time during the experiment, which is in agreement with the findings of Hopkins et al. (2013) in the Arctic but in contrast to Hopkins et al. (2010) and Webb (2015), where iodocarbons were

measured to be significantly lower under elevated $f\text{CO}_2$ (Table 4). Concentrations of all iodocarbons measured in the mesocosms and the Baltic Sea fall within the range of those measured previously in the region (Table 5). Mesocosm concentrations of CH_3I (Fig. 3a) and $\text{C}_2\text{H}_5\text{I}$ (Fig. 3b) showed concentration ranges of 2.91 to 6.25 and 0.23 to 0.76 pmol L^{-1} respectively. CH_3I showed a slight increase in all mesocosms during Phase I, peaking on $t=16$, which corresponded to higher Chl *a* concentrations and correlated throughout the entire experiment with picoeukaryote groups II ($\rho = 0.59$, $p < 0.01$) and III ($\rho = 0.23$, $p < 0.01$; Crawford et al., 2016) and nanoeukaryotes I ($\rho = 0.37$, $p < 0.01$). Significant differences identified between mesocosms for CH_3I were unrelated to elevated $f\text{CO}_2$ ($F = 3.1$, $p < 0.05$), but concentrations were on average 15 % higher in Phase II than Phase I. $\text{C}_2\text{H}_5\text{I}$ decreased slightly during Phases I and II, although concentrations of this halocarbon were close to its detection limit (0.2 pmol L^{-1}), remaining below 1 pmol L^{-1} at all times. As this compound showed no significant effect of elevated $f\text{CO}_2$ and was identified by Orlikowska and Schulz-Bull (2009) as having extremely low concentrations in the Baltic Sea (Table 5), it will not be discussed further.

Table 5. Concentration ranges of trace gases measured in the Baltic Sea compared to concentrations measured in the literature.

Study	DMS concentration range (nmol L ⁻¹)	Halocarbon concentration range (pmol L ⁻¹)						
		CH ₃ I	CH ₂ I ₂	C ₂ H ₅ I	CH ₃ Cl	CHBr ₃	CH ₂ Br ₂	CH ₂ BrCl
SOPRAN Tvärminne Baltic Sea (this study)	1.9–11	4.3–8.6	66.9–374	0.6–1.0	7.0–18	93–192	7.1–10	3.3–5.0
Orlikowska and Schulz-Bull (2009)	0.3–120	1–16	0–85	0.4–1.2	5–50	5.0–40	2.0–10	0.8–2.5
Karlsson et al. (2008)		3.0–7.5				35–60	4.0–7.0	2.0–6.5
Klick and Abrahamsson (1992)			15–709		11–74	14–585		
Klick (1992)			ND–243		ND–57	40–790	ND–86	ND–29
Leck and Rodhe (1991)	0.4–2.8							
Leck et al. (1990)	ND–3.2							

ND – not detected.

No correlation was found between CH₃I and Chl *a* at any phase, and the only correlation of any phytoplankton grouping was with nanoeukaryotes II ($\rho = 0.88$, $p < 0.01$; Crawford et al., 2016). These CH₃I concentrations compare well to the 7.5 pmol L⁻¹ measured by Karlsson et al. (2008) during a cyanobacterial bloom in the Baltic Sea (Table 5) and the summer maximum of 16 pmol L⁻¹ identified by Orlikowska and Schulz-Bull (2009).

Karlsson et al. (2008) showed Baltic Sea halocarbon production occurring predominately during daylight hours, with concentrations at night decreasing by 70 % compared to late afternoon. Light-dependent production of CH₃I has been shown to take place through abiotic processes, including radical recombination of CH₃ and I (Moore and Zafiriou, 1994). However, since samples were integrated over the surface 10 m of the water column, it was impossible to determine whether photochemistry was affecting iodocarbon concentrations near the surface where some UV light was able to pass between the top of the mesocosm film material and the cover. For the same reason, photodegradation of halocarbons (Zika et al., 1984) within the mesocosms was also likely to have been significantly restricted. Thus, as photochemical production was expected to be minimal, biogenic production was likely to have been the dominant source of these compounds. Karlsson et al. (2008) identified *Pseudanabaena* as a key producer of CH₃I in the Baltic Sea. However, the abundance of *Pseudanabaena* was highest during Phase I of the experiment (A. Stühr, personal communication, 2015) when CH₃I concentrations were lower, and as discussed previously, the abundance of these species constituted only a very small proportion of the community. Previous investigations in the laboratory have identified diatoms as significant producers of CH₃I (Hughes et al., 2013; Manley and De La Cuesta, 1997), and the low, steady-state abundance of the diatom populations in the mesocosms could have produced the same relatively steady-state trends in the iodocarbon concentrations.

Measured in the range 57.2–202.2 pmol L⁻¹ in the mesocosms, CH₂I₂ (Fig. 3c) showed the clearest increase in

concentration during Phase II, when it peaked on *t*21 in all mesocosms, with a maximum of 202.2 pmol L⁻¹ in M5 (348 μatm). During Phase II, concentrations of CH₂I₂ were 57 % higher than Phase I and were therefore negatively correlated with Chl *a*. The peak on *t*21 corresponds to the peak identified in DMS on *t*21, and concentrations through all three phases correlate with picoeukaryotes II ($\rho = 0.62$, $p < 0.01$) and III ($\rho = 0.47$, $p < 0.01$) and nanoeukaryotes I ($\rho = 0.88$, $p < 0.01$; Crawford et al., 2015). CH₂ClI (Fig. 3d) showed no peaks during either Phase I or Phase II, remaining within the range of 3.81 to 8.03 pmol L⁻¹ and again correlated with picoeukaryotes groups II ($\rho = 0.34$, $p < 0.01$) and III ($\rho = 0.38$, $p < 0.01$). These results may suggest that these groups possessed halo-peroxidase enzymes able to oxidise I⁻, most likely as an antioxidant mechanism within the cell to remove H₂O₂ (Butler and Carter-Franklin, 2004; Pedersen et al., 1996; Theiler et al., 1978). However, given the lack of response of these compounds to elevated $f\text{CO}_2$ ($F = 1.7$, $p < 0.01$), it is unlikely that production was increased in relation to elevated $f\text{CO}_2$. Production of all iodocarbons increased during Phase II when total Chl *a* decreased, particularly after the walls of the mesocosms were cleaned for the first time, releasing significant volumes of organic aggregates into the water column. Aggregates have been suggested as a source of CH₃I and C₂H₅I (Hughes et al., 2008), likely through the alkylation of inorganic iodide (Urhahn and Ballschmiter, 1998) or through the breakdown of organic matter by microbial activity to supply the precursors required for iodocarbon production (Smith et al., 1992). Hughes et al. (2008) did not identify this route as a pathway for CH₂I₂ or CH₂ClI production, but Carpenter et al. (2005) suggested a production pathway for these compounds through the reaction of HOI with aggregated organic materials.

3.4 Bromocarbons in the mesocosms and the relationships with community composition

No effect of elevated $f\text{CO}_2$ was identified for any of the three bromocarbons, which compared well with the findings from previous mesocosms where bromocarbons were studied

(Hopkins et al., 2010, 2013; Webb, 2015; Table 4). Measured concentrations were comparable to those of Orlikowska and Schulz-Bull (2009) and Karlsson et al. (2008) measured in the southern part of the Baltic Sea (Table 3). The concentrations of CHBr_3 , CH_2Br_2 , and CHBr_2Cl showed no major peaks of production in the mesocosms. CHBr_3 (Fig. 4a) decreased rapidly in all mesocosms over Phase 0 from a maximum measured concentration of $147.5 \text{ pmol L}^{-1}$ in M1 (mean of $138.3 \text{ pmol L}^{-1}$ in all mesocosms) to a mean of $85.7 (\pm 8.2 \text{ SD}) \text{ pmol L}^{-1}$ in all mesocosms for the period t_0 to t_{31} (Phases I and II). The steady-state CHBr_3 concentrations indicated a production source; however, there was no clear correlation with any measured algal groups. CH_2Br_2 concentrations (Fig. 4b) decreased steadily in all mesocosms from $t - 3$ through to t_{31} , over the range 4.0 to 7.7 pmol L^{-1} , and CHBr_2Cl followed a similar trend in the range 1.7 to 4.7 pmol L^{-1} (Fig. 4c). Of the three bromocarbons, only CH_2Br_2 showed correlation with total Chl a ($\rho = 0.52$, $p < 0.01$) and with cryptophyte ($\rho = 0.86$, $p < 0.01$) and dinoflagellate ($\rho = 0.65$, $p < 0.01$)-derived Chl a . Concentrations of CH_2BrI were below detection limit for the entire experiment.

CH_2Br_2 showed positive correlation with Chl a ($\rho = 0.52$, $p < 0.01$), nanoeukaryotes II ($\rho = 0.34$, $p < 0.01$), and cryptophytes ($\rho = 0.86$, $p < 0.01$; see Supplement), whereas CHBr_3 and CHBr_2Cl showed very weak or no correlation with any indicators of algal biomass. Schall et al. (1997) have proposed that CHBr_2Cl is produced in seawater by the nucleophilic substitution of bromide by chloride in CHBr_3 , which given the steady-state concentrations of CHBr_3 would explain the similar distribution of CHBr_2Cl concentrations. Production of all three bromocarbons was identified from large-size cyanobacteria such as *Aphanizomenon flos-aquae* by Karlsson et al. (2008), and in addition, significant correlations were found in the Arabian Sea between the abundance of the cyanobacterium *Trichodesmium* and several bromocarbons (Roy et al., 2011), and the low abundance of such bacteria in the mesocosms would explain the low variation in bromocarbon concentrations through the experiment.

Halocarbon loss processes such as nucleophilic substitution (Moore, 2006), hydrolysis (Elliott and Rowland, 1995), sea–air exchange, and microbial degradation are suggested as of greater importance than the production of these compounds by specific algal groups, particularly given the relatively low growth rates and low net increase in total Chl a . Hughes et al. (2013) identified bacterial inhibition of CHBr_3 production in laboratory cultures of *Thalassiosira* diatoms but that it was not subject to bacterial breakdown, which could explain the relative steady state of CHBr_3 concentrations in the mesocosms. In contrast, significant bacterial degradation of CH_2Br_2 in the same experiments could explain the steady decrease in CH_2Br_2 concentrations seen in the mesocosms. Bacterial oxidation was also identified by Goodwin et al. (1998) as a significant sink for CH_2Br_2 . As discussed for the iodocarbons, photolysis was unlikely due




Figure 4. Mean concentrations (pmol L^{-1}) of (a) CHBr_3 , (b) CH_2Br_2 , and (c) CHBr_2Cl taken from a water sample integrated from the surface 10 m. Dashed lines indicate the phases of the experiment as defined in Fig. 2; $f\text{CO}_2$ shown in the legend is mean $f\text{CO}_2$ across the duration of the experiment.

to the UV absorption of the mesocosm film and limited UV exposure of the surface waters within the mesocosm due to the mesocosm cover. The ratio of CH_2Br_2 to CHBr_3 was also unaffected by increased $f\text{CO}_2$, staying within the range 0.04 to 0.08. This range in ratios is consistent with that calculated by Hughes et al. (2009) in the surface waters of an Antarctic depth profile and attributed to higher sea–air flux of CHBr_3 than CH_2Br_2 due to a greater concentrations gradient, despite the similar transfer velocities of the two compounds (Quack et al., 2007). Using cluster analysis in a time series in the Baltic Sea, Orlikowska and Schulz-Bull (2009) identified

both these compounds as originating from different sources and different pathways of production.

Macroalgal production would not have influenced the mesocosm concentrations after the bags were sealed due to the isolation from the coastal environment. However, macroalgal production into the water column prior to mesocosm installation (Klick, 1992; Leedham et al., 2013; Moore and Tokarczyk, 1993) could account for the high initial concentrations with concentrations decreasing through the duration of the experiment via turnover and transfer to the atmosphere.

3.5 Natural variations in Baltic Sea $f\text{CO}_2$ and the effect on biogenic trace gases

3.5.1 Physical variation and community dynamics

Baltic Sea deep waters have high $f\text{CO}_2$ and subsequently lower pH (Schneider et al., 2002), and the influx to the surface waters surrounding the mesocosms resulted in $f\text{CO}_2$ increasing to $725 \mu\text{atm}$ on $t31$, close to the average $f\text{CO}_2$ of the third-highest mesocosm (M6: $868 \mu\text{atm}$). The input of upwelled water into the region midway through the experiment significantly altered the biogeochemical properties of the waters surrounding the mesocosms, and as a result it is inappropriate to directly compare the community structure and trace gas production of the Baltic Sea and the mesocosms. These conditions imply that pelagic communities in the Baltic Sea are regularly exposed to rapid changes in $f\text{CO}_2$ and the associated pH, as well as having communities associated with the elevated $f\text{CO}_2$ conditions. The changes in biological parameters and trace gas concentrations are therefore discussed here separately from the concentrations measured in the mesocosms.

Given the separation of the waters within the mesocosms and the movement of water masses within the Baltic Sea, it is expected that phytoplankton population structure could be significantly different inside the mesocosms compared to the external waters. Chl a followed the pattern of the mesocosms until $t4$, after which concentrations were significantly higher than any mesocosm, peaking at $6.48 \mu\text{g L}^{-1}$ on $t16$, corresponding to the maximum Chl a peak in the mesocosms and the maximum peak of temperature. As upwelled water intruded into the surface waters, the surface Chl a was diluted with low-Chl a deep water: Chl a in the surface 10 m decreased from around $t16$ at the start of the upwelling until $t31$ when concentrations were once again equivalent to those found in the mesocosms at $1.30 \mu\text{g L}^{-1}$. In addition, there was the potential introduction of different algal groups to the surface, but chlorophytes and cryptophytes were the major contributors to the Chl a in the Baltic Sea, as in the mesocosms. Cyanobacteria contributed less than 2% of the total Chl a in the Baltic Sea (Crawford et al., 2016; Paul et al., 2015).

Temporal community dynamics in the Baltic Sea were very different to that in the mesocosms across the experiment, with euglenophytes, chlorophytes, diatoms, and prasinophytes all showing distinct peaks at the start of Phase II, with these same peaks identified in the nanoeukaryotes I and II and picoeukaryotes II (Crawford et al., 2016; Paul et al., 2015; Supplement Figs. S1 and S2). The decrease in the abundance of many groups during Phase II was attributed to the decrease in temperature and dilution with low-abundance deep waters.

3.5.2 DMS in the Baltic Sea

The Baltic Sea samples gave a mean DMS concentration of $4.6 \pm 2.6 \text{ nmol L}^{-1}$ but peaked at 11.2 nmol L^{-1} on $t16$ and were within the range of previous measurements for the region (Table 5). Strong correlations were seen between DMS and Chl a ($\rho = 0.84$, $p < 0.01$), with the ratio of DMS:Chl a at $1.6 (\pm 0.3) \text{ nmol } \mu\text{g}^{-1}$. Other strong correlations were seen with euglenophytes ($\rho = 0.89$, $p < 0.01$), dinoflagellates ($\rho = 0.61$, $p < 0.05$), and nanoeukaryotes II ($\rho = 0.88$, $p < 0.01$), but no correlation was found between DMS and cyanobacterial abundance or with picoeukaryotes III, which were identified in the mesocosms, suggesting that DMS had a different origin in the Baltic Sea community than in the mesocosms. In addition, the community demands of sulfur are likely to be very different in the Baltic Sea compared to the mesocosms, due to differences in community composition and sulfur availability, and therefore direct comparisons with mesocosm concentrations are inappropriate.

As CO_2 levels increased after $t16$, the DMS concentration measured in the Baltic Sea decreased, from the peak on $t16$ to the lowest recorded sample of the entire experiment at 1.85 nmol L^{-1} on $t31$. As with Chl a , DMS concentrations in the surface of the Baltic Sea may have been diluted with low-DMS deep water

3.5.3 Halocarbon concentrations in the Baltic Sea

Outside the mesocosms in the Baltic Sea, CH_3I was measured at a maximum concentration of 8.65 pmol L^{-1} , during Phase II, and showed a limited effect of the upwelling event. Both CH_2I_2 and CH_2CII showed higher concentrations in the Baltic Sea samples than the mesocosms (CH_2I_2 : $373.9 \text{ pmol L}^{-1}$; CH_2CII : 18.1 pmol L^{-1}) and were correlated with the euglenophytes (CH_2I_2 : $\rho = 0.63$, $p < 0.05$; CH_2CII : $\rho = 0.68$, $p < 0.01$) and nanoeukaryotes II (CH_2I_2 : $\rho = 0.53$, $p < 0.01$; CH_2CII : $\rho = 0.58$, $p < 0.01$), but there was no correlation with Chl a . Both polyhalogenated compounds showed correlation with picoeukaryote groups II and III, indicating that production was probably not limited to a single source. These concentrations of CH_2I_2 and CH_2CII compared well to those measured over a macroalgal bed in the higher-saline waters of the Kattegat by Klick and Abrahamsson (1992), suggesting that macroalgae were a signifi-

cant iodocarbon source in the Baltic Sea. Macroalgal production in the Baltic Sea is likely the predominant iodocarbon source, compared to the mesocosms where macroalgae are excluded.

As with the iodocarbons, the Baltic Sea showed significantly higher concentrations of CHBr_3 ($F = 28.1$, $p < 0.01$), CH_2Br_2 ($F = 208.8$, $p < 0.01$), and CHBr_2Cl ($F = 23.5$, $p < 0.01$) than the mesocosms, with maximum concentrations of 191.6, 10.0, and 5.0 pmol L^{-1} respectively. In the Baltic Sea, only CHBr_3 was correlated with Chl a ($\rho = 0.65$, $p < 0.05$), cyanobacteria ($\rho = 0.61$, $p < 0.01$; Paul et al., 2015), and nanoeukaryotes II ($\rho = 0.56$, $p < 0.01$; Crawford et al., 2016), with the other two bromocarbons showing little to no correlations with any parameter of community activity. Production of bromocarbons from macroalgal sources (Laternus et al., 2000; Leedham et al., 2013; Manley et al., 1992) was likely a significant contributor to the concentrations detected in the Baltic Sea; over the macroalgal beds in the Kattegat, Klick (1992) measured concentrations an order of magnitude higher than seen in this experiment for CH_2Br_2 and CHBr_2Cl . There was only a slight increase in bromocarbon concentrations as a result of the upwelling, indicating that the upwelled water had similar concentrations to the surface waters. These data from the Baltic Sea are presented as an important time series of halocarbon measurements during the summer of 2012 and are expected to add to existing Baltic Sea trace gas datasets.

4 The Baltic Sea as a natural analogue to future ocean acidification?

Mesocosm experiments are a highly valuable tool in assessing the potential impacts of elevated CO_2 on complex marine communities; however, they are limited in that the rapid change in $f\text{CO}_2$ experienced by the community may not be representative of changes in the future ocean (Passow and Riebesell, 2005). This inherent problem with mesocosm experiments can be overcome through using naturally low-pH-high- CO_2 areas such as upwelling regions or vent sites (Hall-Spencer et al., 2008), which can give an insight into populations already living and acclimated to high- CO_2 regimes by exposure over timescales measured in years. This mesocosm experiment was performed at such a location with a relatively high- $f\text{CO}_2$ excursion, which was, however, still low compared to some sites ($800 \mu\text{atm}$ compared to $> 2000 \mu\text{atm}$; Hall-Spencer et al., 2008), and it was clear through the minimal variation in Chl a between all mesocosms that the community was relatively unaffected by elevated $f\text{CO}_2$, although variation could be identified in some phytoplankton groups and some shifts in community composition. The upwelling event occurring midway through our experiment allowed the comparison of the mesocosm findings with a natural analogue of the system, as well as showing the extent to which the system perturbation can occur (up to $800 \mu\text{atm}$).

This event was a fortuitous occurrence during this mesocosm experiment, but as the scale and timing of these upwelling events is difficult to determine, these upwelling events are extremely challenging to study as natural high- CO_2 analogues.

In this paper, we described the temporal changes in concentrations of DMS and halocarbons in natural Baltic phytoplankton communities exposed to elevated- $f\text{CO}_2$ treatments. In contrast to the halocarbons, concentrations of DMS were significantly lower in the highest- $f\text{CO}_2$ treatments compared to the control. Despite very different physicochemical and biological characteristics of the Baltic Sea (e.g. salinity, community composition, and nutrient concentrations), this is a very similar outcome to that seen in several other high- $f\text{CO}_2$ experiments. The Baltic Sea trace gas samples give a good record of trace gas cycling during the injection of high- $f\text{CO}_2$ deep water into the surface community during upwelling events. For the concentrations of halocarbons, the measured concentrations did not change during the upwelling event in the Baltic Sea, which may indicate that emissions of organic iodine and bromine are unlikely to change with future acidification of the Baltic Sea without significant alteration to the meteorological conditions. Further studies of these compounds are important to determine rates of production and consumption to include them in prognostic and predictive models. However, net production of organic sulfur within the Baltic Sea region is likely to decrease with an acidified future ocean scenario, despite the possible acclimation of the microbial community to elevated $f\text{CO}_2$. This will potentially impact the flux of DMS to the atmosphere over northern Europe and could have significant impacts on the local climate through the reduction of atmospheric sulfur aerosols. Data from a previous mesocosm experiment has been used to estimate future global changes in DMS production and predicted that global warming would be amplified (Six et al., 2013); utilising the data from this experiment combined with those of other mesocosm, field, and laboratory experiments and associated modelling provides the basis for a better understanding of the future changes in global DMS production and their climatic impacts.

5 Data availability

Trace gas concentration data are available online from the PANGAEA Data Publisher for Earth and Environmental Science (Webb, 2016; <https://doi.pangaea.de/10.1594/PANGAEA.863649>).

All data regarding carbonate chemistry and chlorophyll a concentrations are available at <https://doi.pangaea.de/10.1594/PANGAEA.86303>. Data for environmental parameters (temperature and salinity) are available at <https://doi.pangaea.de/10.1594/PANGAEA.863116>.

The Supplement related to this article is available online at [doi:10.5194/bg-13-4595-2016-supplement](https://doi.org/10.5194/bg-13-4595-2016-supplement).

Acknowledgements. The Tvärminne 2012 mesocosm experiment was part of the SOPRAN II (Surface Ocean Processes in the Anthropocene) Programme (FKZ 03F0611) and BIOACID II (Biological Impacts of Ocean Acidification) project (FKZ 03F06550), funded by the German Ministry for Education and Research (BMBF) and led by the GEOMAR Helmholtz Centre for Ocean Research Kiel, Germany. The authors thank all participants in the SOPRAN Tvärminne experiment for their assistance, including A. Ludwig for logistical support, the diving team, and the staff of Tvärminne Zoological Research Station for hosting the experiment. We also acknowledge the captain and crew of R/V *ALKOR* (AL394 and AL397) for their work transporting, deploying, and recovering the mesocosms.

This work was funded by a UK Natural Environment Research Council Directed Research Studentship (NE/H025588/1) through the UK Ocean Acidification Research Programme, with CASE funding from Plymouth Marine Laboratory. Additional funding was supplied by the EU Seventh Framework Program (FP7/2007-2013) MESOAQUA (EC Contract no. 228224).

Edited by: H.-P. Grossart

Reviewed by: three anonymous referees

References

- Archer, S. D., Cummings, D., Llewellyn, C., and Fishwick, J.: Phytoplankton taxa, irradiance and nutrient availability determine the seasonal cycle of DMSP in temperate shelf seas, *Mar. Ecol. Prog. Ser.*, 394, 111–124, 2009.
- Archer, S. D., Kimmance, S. A., Stephens, J. A., Hopkins, F. E., Bellerby, R. G. J., Schulz, K. G., Piontek, J., and Engel, A.: Contrasting responses of DMS and DMSP to ocean acidification in Arctic waters, *Biogeosciences*, 10, 1893–1908, doi:10.5194/bg-10-1893-2013, 2013.
- Arnold, H. E., Kerrison, P., and Steinke, M.: Interacting effects of ocean acidification and warming on growth and DMS-production in the haptophyte coccolithophore *Emiliana huxleyi*, *Glob. Change Biol.*, 19, 1007–1016, 2013.
- Avgoustidi, V., Nightingale, P. D., Joint, I., Steinke, M., Turner, S. M., Hopkins, F. E., and Liss, P. S.: Decreased marine dimethyl sulfide production under elevated CO_2 levels in mesocosm and in vitro studies, *Environ. Chem.*, 9, 399–404, 2012.
- Barlow, R. G., Cummings, D. G., and Gibb, S. W.: Improved resolution of mono- and divinyl chlorophylls a and b and zeaxanthin and lutein in phytoplankton extracts using reverse phase C-8 HPLC, *Mar. Ecol. Prog. Ser.*, 161, 303–307, 1997.
- Bentley, R. and Chasteen, T. G.: Environmental VOCs – formation and degradation of dimethyl sulfide, methanethiol and related materials, *Chemosphere*, 55, 291–317, 2004.
- Brussaard, C. P. D., Noordeloos, A. A. M., Witte, H., Collenteur, M. C. J., Schulz, K., Ludwig, A., and Riebesell, U.: Arctic microbial community dynamics influenced by elevated CO_2 levels, *Biogeosciences*, 10, 719–731, doi:10.5194/bg-10-719-2013, 2013.
- Brutemark, A., Engström-Öst, J., and Vehmaa, A.: Long-term monitoring data reveal pH dynamics, trends and variability in the western Gulf of Finland, *Oceanol. Hydrobiol. Stud.*, 40, 91–94, 2011.
- Butler, A. and Carter-Franklin, J. N.: The role of vanadium bromoperoxidase in the biosynthesis of halogenated marine natural products, *Nat. Prod. Rep.*, 21, 180–188, 2004.
- Canadell, J. G., Le Quééré, C., Raupach, M. R., Field, C. B., Buitenhuis, E. T., Ciais, P., Conway, T. J., Gillett, N. P., Houghton, R. A., and Marland, G.: Contributions to accelerating atmospheric CO_2 growth from economic activity, carbon intensity, and efficiency of natural sinks, *P. Natl. Acad. Sci. USA*, 104, 18866–18870, 2007.
- Carpenter, L. J., Hopkins, J. R., Jones, C. E., Lewis, A. C., Parthipan, R., Wevill, D. J., Poissant, L., Pilote, M., and Constant, P.: Abiotic source of reactive organic halogens in the sub-arctic atmosphere?, *Environ. Sci. Technol.*, 39, 8812–8816, 2005.
- Carrión, O., Curson, A. R. J., Kumaresan, D., Fu, Y., Lang, A. S., Mercadé, E., and Todd, J. D.: A novel pathway producing dimethylsulphide in bacteria is widespread in soil environments, *Nat. Commun.*, 6, 6579, doi:10.1038/ncomms7579, 2015.
- Chance, R., Baker, A. R., Küpper, F. C., Hughes, C., Kloareg, B., and Malin, G.: Release and transformations of inorganic iodine by marine macroalgae, *Estuar. Coast. Shelf Sci.*, 82, 406–414, 2009.
- Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, *Nature*, 326, 655–661, 1987.
- Chuck, A. L., Turner, S. M., and Liss, P. S.: Oceanic distributions and air-sea fluxes of biogenic halocarbons in the open ocean, *J. Geophys. Res.*, 110, C10022, doi:10.1029/2004JC002741, 2005.
- Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichet, T., Frielingstein, P., Gao, X., Gutowski, W. J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A. J., and Wehner, M.: Long-term climate change: projections, commitments and irreversibility, in: *Climate Change 2013: The Physical Science Basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M.: Cambridge University Press, Cambridge, UK, available from: <http://dial.academielouvain.be/handle/boreal:140396> (last access: 12 June 2014), 2013.
- Crawford, K. J., Brussaard, C. P. D., and Riebesell, U.: Shifts in the microbial community in the Baltic Sea with increasing CO_2 , *Biogeosciences Discuss.*, doi:10.5194/bg-2015-606, in review, 2016.
- Cubasch, U., Wuebbles, D., Chen, D., Facchini, M. C., Frame, D., Mahowald, N., and Winther, J.-G.: Introduction, in: *Climate Change 2013: The Physical Science Basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK, 2013.
- Curran, M. A. J., Jones, G. B., and Burton, H.: Spatial distribution of dimethylsulfide and dimethylsulfoniopropionate in the Australasian sector of the Southern Ocean, *J. Geophys. Res.*, 103, 16677–16689, 1998.
- Curson, A. R. J., Todd, J. D., Sullivan, M. J., and Johnston, A. W. B.: Catabolism of dimethylsulphoniopropionate: microorganisms, enzymes and genes, *Nat. Rev. Microbiol.*, 9, 849–859, 2011.

- Czerny, J., Schulz, K. G., Krug, S. A., Ludwig, A., and Riebesell, U.: Technical Note: The determination of enclosed water volume in large flexible-wall mesocosms “KOSMOS”, *Biogeosciences*, 10, 1937–1941, doi:10.5194/bg-10-1937-2013, 2013.
- del Valle, D. A., Slezak, D., Smith, C. M., Rellinger, A. N., Kieber, D. J., and Kiene, R. P.: Effect of acidification on preservation of DMSP in seawater and phytoplankton cultures: Evidence for rapid loss and cleavage of DMSP in samples containing *Phaeocystis* sp., *Mar. Chem.*, 124, 57–67, 2011.
- Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean acidification: the other CO_2 problem, *Annu. Rev. Mar. Sci.*, 1, 169–192, 2009.
- Drotar, A., Burton, G. A., Tavernier, J. E., and Fall, R.: Widespread occurrence of bacterial thiol methyltransferases and the biogenic emission of methylated sulfur gases, *Appl. Environ. Microbiol.*, 53, 1626–1631, 1987.
- Elliott, S. and Rowland, F. S.: Methyl halide hydrolysis rates in natural waters, *J. Atmos. Chem.*, 20, 229–236, 1995.
- Endres, S., Galgani, L., Riebesell, U., Schulz, K.-G., and Engel, A.: Stimulated bacterial growth under elevated $p\text{CO}_2$: results from an off-shore mesocosm study, *PLoS One*, 9, e99228, doi:10.1371/journal.pone.0099228, 2014.
- Engel, A., Schulz, K. G., Riebesell, U., Bellerby, R. G. J., Delille, B., and Schartau, M.: Effects of CO_2 on particle size distribution and phytoplankton abundance during a mesocosm bloom experiment (PeECE II), *Biogeosciences*, 5, 509–521, doi:10.5194/bg-5-509-2008, 2008.
- Engel, A., Borchard, C., Piontek, J., Schulz, K. G., Riebesell, U., and Bellerby, R. G. J.: CO_2 increases ^{14}C primary production in an Arctic plankton community, *Biogeosciences*, 10, 1291–1308, doi:10.5194/bg-10-1291-2013, 2013.
- Finni, T., Kononen, K., Olsonen, R., and Wallström, K.: The History of Cyanobacterial Blooms in the Baltic Sea, *AMBIO A J. Hum. Environ.*, 30, 172–178, 2001.
- Franklin, D. J., Poulton, A. J., Steinke, M., Young, J., Peeken, I., and Malin, G.: Dimethylsulphide, DMSP-lyase activity and microplankton community structure inside and outside of the Mauritanian upwelling, *Prog. Oceanogr.*, 83, 134–142, 2009.
- Gidhagen, L.: Coastal upwelling in the Baltic Sea – Satellite and in situ measurements of sea-surface temperatures indicating coastal upwelling, *Estuar. Coast. Shelf Sci.*, 24, 449–462, 1987.
- Goodwin, K., Schaefer, J. K., and Oremland, R. S.: Bacterial oxidation of dibromomethane and methyl bromide in natural waters and enrichment cultures, *Appl. Environ. Microbiol.*, 64, 4629–4636, 1998.
- Goodwin, K. D., North, W. J., and Lidstrom, M. E.: Production of bromoform and dibromomethane by giant kelp: factors affecting release and comparison to anthropogenic bromine sources, *Limnol. Oceanogr.*, 42, 1725–1734, 1997.
- Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S. M., Rowley, S. J., Tedesco, D., and Buia, M.-C.: Volcanic carbon dioxide vents show ecosystem effects of ocean acidification, *Nature*, 454, 96–99, 2008.
- Hartmann, D. L., Klein Tank, A. M. G., Rusticucci, M., Alexander, L. V., Bronnimann, S., Charabi, Y., Dentener, F. J., Dlugokencky, E. J., Easterling, D. R., Kaplan, A., Soden, B. J., Thorne, P. W., Wild, M., and Zhai, P. M.: Observations: Atmosphere and Surface, in: *Climate Change 2013: The Physical Science Basis*, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, Cambridge, UK, 2013.
- Hatton, A. D., Darroch, L., and Malin, G.: The role of dimethylsulphoxide in the marine biogeochemical cycle of dimethylsulphide, *Oceanogr. Mar. Biol. Annu. Rev.*, 42, 29–56, 2004.
- Hofmann, G. E., Barry, J. P., Edmunds, P. J., Gates, R. D., Hutchins, D. A., Klinger, T., and Sewell, M. A.: The effect of ocean acidification on calcifying organisms in marine ecosystems: an organism-to-ecosystem perspective, *Annu. Rev. Ecol. Evol. Syst.*, 41, 127–147, 2010.
- Hopkins, F. E. and Archer, S. D.: Consistent increase in dimethyl sulfide (DMS) in response to high CO_2 in five shipboard bioassays from contrasting NW European waters, *Biogeosciences*, 11, 4925–4940, doi:10.5194/bg-11-4925-2014, 2014.
- Hopkins, F. E., Turner, S. M., Nightingale, P. D., Steinke, M., Bakker, D., and Liss, P. S.: Ocean acidification and marine trace gas emissions, *P. Natl. Acad. Sci. USA*, 107, 760–765, 2010.
- Hopkins, F. E., Kimmance, S. A., Stephens, J. A., Bellerby, R. G. J., Brussaard, C. P. D., Czerny, J., Schulz, K. G., and Archer, S. D.: Response of halocarbons to ocean acidification in the Arctic, *Biogeosciences*, 10, 2331–2345, doi:10.5194/bg-10-2331-2013, 2013.
- Hornick, T., Bach, L. T., Crawford, K. J., Spilling, K., Achterberg, E. P., Brussaard, C. P. D., Riebesell, U., and Grossart, H.-P.: Ocean acidification indirectly alters trophic interaction of heterotrophic bacteria at low nutrient conditions, *Biogeosciences Discuss.*, doi:10.5194/bg-2016-61, in review, 2016.
- Hughes, C., Malin, G., Nightingale, P. D., and Liss, P. S.: The effect of light stress on the release of volatile iodocarbons by three species of marine microalgae, *Limnol. Oceanogr.*, 51, 2849–2854, 2006.
- Hughes, C., Malin, G., Turley, C. M., Keely, B. J., Nightingale, P. D., and Liss, P. S.: The production of volatile iodocarbons by biogenic marine aggregates, *Limnol. Oceanogr.*, 53, 867–872, 2008.
- Hughes, C., Chuck, A. L., Rossetti, H., Mann, P. J., Turner, S. M., Clarke, A., Chance, R., and Liss, P. S.: Seasonal cycle of seawater bromoform and dibromomethane concentrations in a coastal bay on the western Antarctic Peninsula, *Global Biogeochem. Cy.*, 23, GB2024, doi:10.1029/2008GB003268, 2009.
- Hughes, C., Franklin, D. J., and Malin, G.: Iodomethane production by two important marine cyanobacteria: *Prochlorococcus marinus* (CCMP 2389) and *Synechococcus* sp. (CCMP 2370), *Mar. Chem.*, 125, 19–25, 2011.
- Hughes, C., Johnson, M., Utting, R., Turner, S., Malin, G., Clarke, A., and Liss, P. S.: Microbial control of bromocarbon concentrations in coastal waters of the western Antarctic Peninsula, *Mar. Chem.*, 151, 35–46, 2013.
- Janssen, F., Schrum, C., and Backhaus, J.: A climatological data set of temperature and salinity for the Baltic Sea and the North Sea, *Dtsch. Hydrogr. Zeitschrift*, Supplement available from: <http://link.springer.com/article/10.1007/BF02933676> (last access: 16 August 2014), 1999.
- Kangro, K., Olli, K., Tamminen, T., and Lignell, R.: Species-specific responses of a cyanobacteria-dominated phytoplankton community to artificial nutrient limitation in the Baltic Sea, *Mar. Ecol. Prog. Ser.*, 336, 15–27, 2007.

- Karlsson, A., Auer, N., Schulz-Bull, D., and Abrahamsson, K.: Cyanobacterial blooms in the Baltic – A source of halocarbons, *Mar. Chem.*, 110, 129–139, 2008.
- Kiene, R. P. and Hines, M. E.: Microbial formation of dimethyl sulfide in anoxic sphagnum peat, *Appl. Environ. Microbiol.*, 61, 2720–2726, 1995.
- Kiene, R. P. and Linn, L. J.: Distribution and turnover of dissolved DMSP and its relationship with bacterial production and dimethylsulfide in the Gulf of Mexico, *Limnol. Oceanogr.*, 45, 849–861, 2000.
- Kiene, R. P. and Visscher, P. T.: Production and fate of methylated sulfur compounds from methionine and dimethylsulfoniopropionate in anoxic salt marsh sediments, *Appl. Environ. Microbiol.*, 53, 2426–2434, 1987.
- Kim, J.-M., Lee, K., Yang, E. J., Shin, K., Noh, J. H., Park, K.-T., Hyun, B., Jeong, H.-J., Kim, J.-H., Kim, K. Y., Kim, M., Kim, H.-C., Jang, P.-G., and Jang, M.-C.: Enhanced production of oceanic dimethylsulfide resulting from CO_2 -induced grazing activity in a high CO_2 world., *Environ. Sci. Technol.*, 44, 8140–8143, 2010.
- Klick, S.: Seasonal variations of biogenic and anthropogenic halocarbons in seawater from a coastal site, *Limnol. Oceanogr.*, 37, 1579–1585, 1992.
- Klick, S. and Abrahamsson, K.: Biogenic volatile iodated hydrocarbons in the ocean, *J. Geophys. Res.*, 97, 12683–12687, 1992.
- Lana, A., Simó, R., Vallina, S. M., and Dachs, J.: Re-examination of global emerging patterns of ocean DMS concentration, *Biogeochemistry*, 110, 173–182, 2012.
- Laturmus, F., Giese, B., Wiencke, C., and Adams, F. C.: Low-molecular-weight organoiodine and organobromine compounds released by polar macroalgae—the influence of abiotic factors, *Fresenius. J. Anal. Chem.*, 368, 297–302 2000.
- Leck, C. and Rodhe, H.: Emissions of marine biogenic sulfur to the atmosphere of northern Europe, *J. Atmos. Chem.*, 12, 63–86, 1991.
- Leck, C., Larsson, U., Bågander, L. E., Johansson, S., and Hajdu, S.: Dimethyl sulfide in the Baltic Sea: annual variability in relation to biological activity, *J. Geophys. Res.*, 95, 3353–3363, 1990.
- Lee, G., Park, J., Jang, Y., Lee, M., Kim, K. R., Oh, J. R., Kim, D., Yi, H. I., and Kim, T. Y.: Vertical variability of seawater DMS in the South Pacific Ocean and its implication for atmospheric and surface seawater DMS, *Chemosphere*, 78, 1063–1070, 2010.
- Leedham, E. C., Hughes, C., Keng, F. S. L., Phang, S.-M., Malin, G., and Sturges, W. T.: Emission of atmospherically significant halocarbons by naturally occurring and farmed tropical macroalgae, *Biogeosciences*, 10, 3615–3633, doi:10.5194/bg-10-3615-2013, 2013.
- Lehmann, A. and Myrberg, K.: Upwelling in the Baltic Sea – A review, *J. Mar. Syst.*, 74, S3–S12, 2008.
- Lischka, S., Bach, L. T., Schulz, K.-G., and Riebesell, U.: Micro- and mesozooplankton community response to increasing CO_2 levels in the Baltic Sea: insights from a large-scale mesocosm experiment, *Biogeosciences Discuss.*, 12, 20025–20070, doi:10.5194/bgd-12-20025-2015, 2015.
- Liss, P., Marandino, C. A., Dahl, E., Helmig, D., Hints, E. J., Hughes, C., Johnson, M., Moore, R. M., Plane, J. M. C., Quack, B., Singh, H. B., Stefels, J., von Glasow, R., and Williams, J.: Short-lived trace gases in the surface ocean and the atmosphere, in: *Ocean-Atmosphere Interactions of Gases and Particles*, edited by: Liss, P. and Johnson, M., 55–112, 2014.
- Lomans, B. P., Smolders, A., Intven, L. M., Pol, A., Op, D., and van der Drift, C.: Formation of dimethyl sulfide and methanethiol in anoxic freshwater sediments, *Appl. Environ. Microbiol.*, 63, 4741–4747, 1997.
- Mackey, M. D., Mackey, D. J., Higgins, H. W., and Wright, S. W.: CHEMTAX a program for estimating class abundances from chemical markers: application to HPLC measurements of phytoplankton, *Mar. Ecol. Prog. Ser.*, 144, 265–283, 1996.
- Manley, S. L. and De La Cuesta, J. L.: Methyl iodide production from marine phytoplankton cultures, *Limnol. Oceanogr.*, 42, 142–147, 1997.
- Manley, S. L., Goodwin, K., and North, W. J.: Laboratory production of bromoform, methylene bromide, and methyl iodide by macroalgae in and distribution nearshore Southern California waters, *Limnol. Oceanogr.*, 37, 1652–1659, 1992.
- Martino, M., Liss, P. S., and Plane, J. M. C.: The photolysis of dihalomethanes in surface seawater, *Environ. Sci. Technol.*, 39, 7097–7101 2005.
- Moore, R. M.: Methyl halide production and loss rates in sea water from field incubation experiments, *Mar. Chem.*, 101, 213–219, 2006.
- Moore, R. M. and Tokarczyk, R.: Volatile biogenic halocarbons in the northwest Atlantic, *Global Biogeochem. Cy.*, 7, 195–210, 1993.
- Moore, R. M. and Zafiriou, O. C.: Photochemical production of methyl iodide in seawater, *J. Geophys. Res.*, 99, 16415–16420, 1994.
- National Measurement Institute of Australia: Proficiency Study 12–23: DMS in seawater, 2013.
- Nausch, M., Nausch, G., Lass, H. U., Mohrholz, V., Nagel, K., Siegel, H., and Wasmund, N.: Phosphorus input by upwelling in the eastern Gotland Basin (Baltic Sea) in summer and its effects on filamentous cyanobacteria, *Estuar. Coast. Shelf Sci.*, 83, 434–442, 2009.
- Newbold, L. K., Oliver, A. E., Booth, T., Tiwari, B., DeSantis, T., Maguire, M., Andersen, G., van der Gast, C. J., and Whiteley, A. S.: The response of marine picoplankton to ocean acidification, *Environ. Microbiol.*, 14, 2293–2307, 2012.
- Niemisto, L., Rinne, I., and Melvasalo, T.: Blue-green algae and their nitrogen fixation in the Baltic Sea in 1980, 1982 and 1984, *Meri*, 17, 1–59, 1989.
- Oberg, J.: Cyanobacterial blooms in the Baltic Sea in 2013, *HELCOM Balt, Sea Environ, Fact Sheet*, 2013.
- O’Dowd, C. D., Jimenez, J. L., Bahreini, R., Flagan, R. C., Seinfeld, J. H., Hameri, K., Pirjola, L., Kulmala, M., Jennings, S. G., and Hoffmann, T.: Marine aerosol formation from biogenic iodine emissions, *Nature*, 417, 632–636, 2002.
- Orlikowska, A. and Schulz-Bull, D. E.: Seasonal variations of volatile organic compounds in the coastal Baltic Sea, *Environ. Chem.*, 6, 495–507, 2009.
- Park, K.-T., Lee, K., Shin, K., Yang, E. J., Hyun, B., Kim, J.-M., Noh, J. H., Kim, M., Kong, B., Choi, D. H., Choi, S.-J., Jang, P.-G., and Jeong, H. J.: Direct linkage between dimethyl sulfide production and microzooplankton grazing, resulting from prey composition change under high partial pressure of carbon dioxide conditions, *Environ. Sci. Technol.*, 48, 4750–4756, 2014.
- Passow, U. and Riebesell, U.: Mesocosm perturbation experiments and the sensitivity of marine biological systems to global change, *Solas News*, 1, 12–13, 2005.

- Paul, A. J., Bach, L. T., Schulz, K.-G., Boxhammer, T., Czerny, J., Achterberg, E. P., Hellemann, D., Trense, Y., Nausch, M., Sswat, M., and Riebesell, U.: Effect of elevated CO_2 on organic matter pools and fluxes in a summer Baltic Sea plankton community, *Biogeosciences*, 12, 6181–6203, doi:10.5194/bg-12-6181-2015, 2015.
- Paul, A., Schulz, K. G., Achterberg, E., Hellemann, D., Nausch, M., Boxhammer, T., Bach, L., and Trense, Y.: KOSMOS Finland 2012 mesocosm study: carbonate chemistry, particulate and dissolved matter pools, and phytoplankton community composition using marker pigments (CHEMTAX), <https://doi.pangaea.de/10.1594/PANGAEA.863032>, 2016.
- Pedersen, M., Collen, J., Abrahamsson, K., and Ekdahl, A.: Production of halocarbons from seaweeds: an oxidative stress reaction?, *Sci. Mar.*, 60 (Supplement 1), 257–263, 1996.
- Piontek, J., Borchard, C., Sperling, M., Schulz, K. G., Riebesell, U., and Engel, A.: Response of bacterioplankton activity in an Arctic fjord system to elevated $p\text{CO}_2$: results from a mesocosm perturbation study, *Biogeosciences*, 10, 297–314, doi:10.5194/bg-10-297-2013, 2013.
- Quack, B., Peeken, I., Petrick, G., and Nachtigall, K.: Oceanic distribution and sources of bromoform and dibromomethane in the Mauritanian upwelling, *J. Geophys. Res.*, 112, C10006, doi:10.1029/2006JC003803, 2007.
- Quinn, P. K. and Bates, T. S.: The case against climate regulation via oceanic phytoplankton sulphur emissions, *Nature*, 480, 51–56, 2011.
- Raateoja, M., Kuosa, H., and Hällfors, S.: Fate of excess phosphorus in the Baltic Sea: A real driving force for cyanobacterial blooms?, *J. Sea Res.*, 65, 315–321, 2011.
- Raven, J. R., Caldeira, K., Elderfield, H., Hoegh-Guldberg, O., Liss, P. S., Riebesell, U., Shepherd, J., Turley, C., and Watson, A.: Ocean acidification due to increasing atmospheric carbon dioxide, *R. Soc. Policy Doc.*, 12, 1–51, 2005.
- Riebesell, U., Czerny, J., von Bröckel, K., Boxhammer, T., Büdenbender, J., Deckelnick, M., Fischer, M., Hoffmann, D., Krug, S. A., Lentz, U., Ludwig, A., Mucche, R., and Schulz, K. G.: Technical Note: A mobile sea-going mesocosm system – new opportunities for ocean change research, *Biogeosciences*, 10, 1835–1847, doi:10.5194/bg-10-1835-2013, 2013.
- Ross, P. M., Parker, L., O'Connor, W. A., and Bailey, E. A.: The impact of ocean acidification on reproduction, early development and settlement of marine organisms, *Water*, 3, 1005–1030, 2011.
- Roy, R., Pratihary, A., Narvenkar, G., Mochemadkar, S., Gauns, M., and Naqvi, S. W. A.: The relationship between volatile halocarbons and phytoplankton pigments during a *Trichodesmium* bloom in the coastal eastern Arabian Sea, *Estuar. Coast. Shelf Sci.*, 95, 110–118, 2011.
- Scarratt, M. G. and Moore, R. M.: Production of methyl bromide and methyl chloride in laboratory cultures of marine phytoplankton II, *Mar. Chem.*, 59, 311–320, 1998.
- Schall, C., Heumann, K. G., and Kirst, G. O.: Biogenic volatile organoiodine and organobromine hydrocarbons in the Atlantic Ocean from 42° N to 72° S, *Fresenius. J. Anal. Chem.*, 359, 298–305, 1997.
- Schneider, B., Nausch, G., Kubsch, H., and Petersohn, I.: Accumulation of total CO_2 during stagnation in the Baltic Sea deep water and its relationship to nutrient and oxygen concentrations, *Mar. Chem.*, 77, 277–291, 2002.
- Schulz, K. G., Bellerby, R. G. J., Brussaard, C. P. D., Büdenbender, J., Czerny, J., Engel, A., Fischer, M., Koch-Klavsen, S., Krug, S. A., Lischka, S., Ludwig, A., Meyerhöfer, M., Nondal, G., Silyakova, A., Stühr, A., and Riebesell, U.: Temporal biomass dynamics of an Arctic plankton bloom in response to increasing levels of atmospheric carbon dioxide, *Biogeosciences*, 10, 161–180, doi:10.5194/bg-10-161-2013, 2013.
- Schulz, K. G.: KOSMOS Finland 2012 mesocosm study: environmental parameter, <https://doi.pangaea.de/10.1594/PANGAEA.863116>, 2016.
- Simó, R., Archer, S. D., Pedros-Alio, C., Gilpin, L., and Stelfox-Widdicombe, C. E.: Coupled dynamics of dimethylsulfoniopropionate and dimethylsulfide cycling and the microbial food web in surface waters of the North Atlantic, *Limnol. Oceanogr.*, 47, 53–61, 2002.
- Simó, R., Vila-Costa, M., Alonso-Sáez, L., Cardelús, C., Guadayol, Ó., Vázquez-Dominguez, E., and Gasol, J. M.: Annual DMSP contribution to S and C fluxes through phytoplankton and bacterioplankton in a NW Mediterranean coastal site, *Aquat. Microb. Ecol.*, 57, 43–55, 2009.
- Six, K. D., Kloster, S., Ilyina, T., Archer, S. D., Zhang, K., and Maier-Reimer, E.: Global warming amplified by reduced sulphur fluxes as a result of ocean acidification, *Nature Climatic Change*, 3, 975–978, 2013.
- Smith, D. C., Simon, M., Alldredge, A. L., and Azam, F.: Intense hydrolytic enzyme activity on marine aggregates and implications for rapid particle dissolution, *Nature*, 359, 139–142, 1992.
- Solomon, S., Garcia, R. R., and Ravishankara, A. R.: On the role of iodine in ozone depletion, *J. Geophys. Res.*, 99, 20491–20499, 1994.
- Stal, L. J., Albertano, P., Bergman, B., von Bröckel, K., Gallon, J. R., Hayes, P. K., Sivonen, K., and Walsby, A. E.: BASIC: Baltic Sea cyanobacteria. An investigation of the structure and dynamics of water blooms of cyanobacteria in the Baltic Sea – responses to a changing environment, *Cont. Shelf Res.*, 23, 1695–1714, 2003.
- Stets, E. G., Hines, M. E., and Kiene, R. P.: Thiol methylation potential in anoxic, low-pH wetland sediments and its relationship with dimethylsulfide production and organic carbon cycling, *FEMS Microbiol. Ecol.*, 47, 1–11, 2004.
- Sunda, W., Kieber, D. J., Kiene, R. P., and Huntsman, S.: An antioxidant function for DMSP and DMS in marine algae, *Nature*, 418, 317–320, 2002.
- Theiler, R., Cook, J. C., Hager, L. P., and Siuda, J. F.: Halohydrocarbon synthesis by bromoperoxidase, *Science*, 202, 1094–1096, 1978.
- Turner, S. M., Malin, G., Liss, P. S., Harbour, D. S., and Holligan, P. M.: The seasonal variation of dimethyl sulfide and dimethylsulfoniopropionate concentrations in nearshore waters, *Limnol. Oceanogr.*, 33, 364–375, 1988.
- Urhahn, T. and Ballschmiter, K.: Chemistry of the biosynthesis of halogenated methanes: C1-organohalogens as pre-industrial chemical stressors in the environment?, *Chemosphere*, 37, 1017–1032, 1998.
- Vila-Costa, M., Simó, R., Harada, H., Gasol, J. M., Slezak, D., and Kiene, R. P.: Dimethylsulfoniopropionate uptake by marine phytoplankton, *Science*, 314, 652–4, 2006a.
- Vila-Costa, M., del Valle, D. A., González, J. M., Slezak, D., Kiene, R. P., Sánchez, O., and Simó, R.: Phylogenetic identification and

- metabolism of marine dimethylsulfide-consuming bacteria, *Environ. Microbiol.*, 8, 2189–2200, 2006b.
- Visscher, P. T., Baumgartner, L. K., Buckley, D. H., Rogers, D. R., Hogan, M. E., Raleigh, C. D., Turk, K. A., and Des Marais, D. J.: Dimethyl sulphide and methanethiol formation in microbial mats: potential pathways for biogenic signatures, *Environ. Microbiol.*, 5, 296–308, 2003.
- Vogt, M., Steinke, M., Turner, S. M., Paulino, A., Meyerhöfer, M., Riebesell, U., LeQuéré, C., and Liss, P. S.: Dynamics of dimethylsulphoniopropionate and dimethylsulphide under different CO_2 concentrations during a mesocosm experiment, *Biogeosciences*, 5, 407–419, doi:10.5194/bg-5-407-2008, 2008.
- Wasmund, N.: Occurrence of cyanobacterial blooms in the Baltic Sea in relation to environmental conditions, *Iny. Rev. Ges. Hydrobiol.*, 82, 169–184, 1997.
- Webb, A.: The effects of elevated CO_2 and ocean acidification on the production of marine biogenic trace gases, PhD Thesis, Univ. East Angl., March 2015, 235–299, 2015.
- Webb, A. L., Malin, G., Hopkins, F. E., Ho, K.-L., Riebesell, U., Schulz, K., Larsen, A., and Liss, P.: Ocean acidification has different effects on the production of dimethylsulphide and dimethylsulphoniopropionate measured in cultures of *Emiliania huxleyi* RCC1229 and mesocosm study: a comparison of laboratory monocultures and community interactions, *Environ. Chem.*, 13, EN14268, doi:10.1071/EN14268, 2015.
- Webb, A., Leedham-Elvidge, E., Hughes, C., Hopkins, F., Malin, G., von Glasow, R., and Liss, P.: Dimethylsulphide and halo-carbon concentrations taken from 10 m integrated samples during the 2012 KOSMOS Finland Mesocosm experiment, <https://doi.pangaea.de/10.1594/PANGAEA.863649>, 2016.
- Welschmeyer, N. A.: Fluorometric analysis of chlorophyll *a* in the presence of chlorophyll *b* and pheopigments, *Limnol. Oceanogr.*, 39, 1985–1992, 1994.
- Zika, R. G., Gidel, L. T., and Davis, D. D.: A comparison of photolysis and substitution decomposition rates of methyl iodide in the ocean, *Geophys. Res. Lett.*, 11, 353–356, 1984.
- Zinder, S. H., Doemel, W. N., and Brock, T. D.: Production of volatile sulfur compounds during the decomposition of algal mats, *Appl. Environ. Microbiol.*, 34, 859–861, 1977.
- Zindler, C., Peeken, I., Marandino, C. A., and Bange, H. W.: Environmental control on the variability of DMS and DMSP in the Mauritanian upwelling region, *Biogeosciences*, 9, 1041–1051, doi:10.5194/bg-9-1041-2012, 2012.