

Salinity change in the subtropical Atlantic: Secular increase and teleconnections to the North Atlantic Oscillation

Brad E. Rosenheim,¹ Peter K. Swart,¹ Simon R. Thorrold,² Anton Eisenhauer,³ and Philippe Willenz⁴

Received 14 September 2004; revised 18 October 2004; accepted 6 December 2004; published 21 January 2005.

[1] Recent studies comparing shipboard data between the 1950's and the 1990's have shown significant, heterogeneous adjustments of the temperature-salinity structure of the N. Atlantic Ocean. Here, we present proxy records of temperature and salinity from aragonite sclerosponge skeletons, extending existing records of the Salinity Maximum Waters (SMW) of the N. Atlantic back to 1890. These proxy records show secular temperature increases of 1.6–2.0°C, higher than published global averages, and salinity increases of 0.35–0.5 psu, smaller than short-term secular trends recently measured. Salinity reconstructions vary more significantly on the decadal scale, showing changes that are related to low-frequency variations of the North Atlantic Oscillation (NAO). On both secular and decadal time scales, the records indicate significant thermohaline changes in the SMW, either via forcing at the surface or increasing depths of density surfaces in the Bahamas. **Citation:** Rosenheim, B. E., P. K. Swart, S. R. Thorrold, A. Eisenhauer, and P. Willenz (2005), Salinity change in the subtropical Atlantic: Secular increase and teleconnections to the North Atlantic Oscillation, *Geophys. Res. Lett.*, 32, L02603, doi:10.1029/2004GL021499.

1. Introduction

[2] There have been numerous observations that global temperatures have been rising through the last century, generating significant interest in changes of the storage and transfer of heat between ocean and atmosphere [Cardone *et al.*, 1990; Levitus *et al.*, 2000; Barnett *et al.*, 2001; Folland *et al.*, 2001; Levitus *et al.*, 2001]. The broad-reaching effects of increasing global heat content [Folland *et al.*, 2001] as well as oceanic heat content [Levitus *et al.*, 2000, 2001] include wind stress changes [Cardone *et al.*, 1990], salinity increases [Curry *et al.*, 2003], and temperature increases, however instrumental records are limited in their time domain. All of these changes indicate the varied mechanisms of the transfer of heat and momentum across

the tropical air-sea interface. Compared to natural variability in other proxy records stretching as far as 6 centuries, there is evidence that these changes are unprecedented in their global significance [Haase-Schramm *et al.*, 2003; Mann and Jones, 2003].

[3] Salinity changes in the N. Atlantic, including the Caribbean Sea, play an important role in global change because of the significant regional and global transport of the highly saline waters formed in the basin. The salinity and temperature profiles of the diffuse return flow of the N. Atlantic gyre influence the formation of subtropical cells (STC's) [Zhang *et al.*, 2003] and of North Atlantic Deep Water (NADW) [Blanke *et al.*, 2002] which both redistribute heat taken up by the ocean in lower latitudes. The effects of salinity change on the latter include such large-scale climate instability such the deglaciation of the northern hemisphere marking the beginning of the Holocene [Broecker *et al.*, 1989; Broecker, 1994; Schmidt *et al.*, 2004]. More recently, instrumental records have indicated a salinity adjustment of the Atlantic Ocean [Curry *et al.*, 2003], which may be linked to the century long increase of CO₂ emissions associated with industrialization [Levitus *et al.*, 2000, 2001] or a positive excursion of the NAO index, the pressure difference between Iceland and the Açores [Marshall *et al.*, 2001]. Curry *et al.* [2003] present evidence of salinity increase in the subtropics, potentially increasing both the amount of water subducted into the STC's and the salinity of waters delivered back to the gyre system.

[4] In this study, we analyze two continuous century long proxy records of salinity and temperature from sclerosponges inhabiting two of the Bahamas' deep channels (Figure 1). This area represents a crucial junction between Caribbean waters and N. Atlantic Gyre return flow waters from the SMW formed in the eastern Atlantic between 20°N and 25°N when intrusion of hot, arid Saharan air masses result in high net evaporation. The SMW subduct along density surfaces to constitute the subsurface waters of the Bahamas and the interior Caribbean, known as the Subtropical Underwater (SUW). In the Bahamas, subsurface waters adjacent to large shelf areas can also show periodic influence of high-density bank top plume waters that form locally as hypersaline waters cool and are entrained over the deep channels by cold fronts [Hickey *et al.*, 2000].

2. Methods and Approach

[5] Our reconstruction of SMW salinity and temperature is derived from skeletal proxies of two specimens of the sclerosponge *Ceratoporella nicholsoni* taken from water

¹Division of Marine Geology and Geophysics, Rosenstiel School of Marine and Atmospheric Sciences, University of Miami, Miami, Florida, USA.

²Biology Department, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts, USA.

³Department for Marine, Environmental Sciences, Research Center for Marine Geosciences (GEOMAR), Kiel, Germany.

⁴Section Malacology, Department of Invertebrates, CALMARS Group, Royal Belgian Institute of Natural Sciences (IRSNB/KBIN), Brussels, Belgium.




Figure 1. Map of the Bahamas with open circles showing locations of the analyzed sclerosponges. The Bahamas are shown in black and the shallow Bahamas bank waters are outlined in gray. The Exuma sclerosponge was collected from 136 m and the TOTO sclerosponge from 143 m. The sclerosponge collected from TOTO is adjacent to a much larger shelf area than the Exuma sclerosponge.

depths of 136 m (Exuma Sound) and 143 m (Tongue of the Ocean (TOTO); Figure 1). The Atlantic sclerosponge *C. nicholsoni* lives in cryptic environments up to 250 m deep and secretes a dense aragonite skeleton at rates of approximately 200 $\mu\text{m}/\text{y}$ [Lang *et al.*, 1975; Benavides and Druffel, 1986; Willenz and Hartman, 1999]. These organisms deposit their skeletons in isotopic equilibrium with water [Böhm *et al.*, 2000], recording both ambient water temperature and $\delta^{18}\text{O}_{\text{sw}}$, which is related to salinity. A recent in situ calibration of Sr/Ca ratios to temperature for this species [Rosenheim *et al.*, 2004] allows differentiation between the temperature and salinity components of the $\delta^{18}\text{O}_{\text{arag}}$ records of these organisms. The temperature dependence of Sr/Ca ratios in sclerosponges is 2–3 times more sensitive than that of inorganic aragonite, making this calibration less sensitive to the assumption of invariant $\text{Sr}/\text{Ca}_{\text{sw}}$ than studies using zooxanthellate corals.

[6] The sclerosponges were dated using U/Th radiometric techniques and then analyzed using high resolution micro-milling for oxygen isotope analysis via isotope ratio mass spectrometry and laser ablation sampling for Sr/Ca analysis via ICP-MS. Constant-growth age models constructed from U/Th dating indicate growth rates of 145 $\mu\text{m}\cdot\text{y}^{-1}$ for the Exuma sclerosponge and 172 $\mu\text{m}\cdot\text{y}^{-1}$ for the TOTO sclerosponge (auxiliary material¹). The radiometric age model of the TOTO sclerosponge was tuned to annual Sr/Ca variations after Swart *et al.* [2002], however because tuned age models and constant growth models are within 5% [Swart *et al.*, 2002], the U/Th constant growth model was used for the Exuma sclerosponge. Micro-milling and laser

ablation were performed from mirror-image slabs of sclerosponge skeleton prepared from the same thick slab, enabling straightforward correlation between isotope and minor element transects. Measurement methods are summarized and results are shown in the auxiliary material as well as in previous work on these sclerosponges [Swart *et al.*, 2002; Rosenheim *et al.*, 2004].

3. Time Series of Proxy Temperature and Salinity

[7] Greater high-frequency variation is present for both Sr/Ca and $\delta^{18}\text{O}_{\text{arag}}$ in the TOTO sclerosponge (Figure 2) due to higher sampling resolution (0.01 mm vs. 0.02 mm) and the presence of a larger shelf area where dense bank-top plumes can form (Figure 1). Linear trends in $\delta^{18}\text{O}_{\text{arag}}$ and Sr/Ca of both sclerosponges decreased between 1890–1990, indicating temperature increase (Figure 2). Temperature increases indicated by Sr/Ca were 1.6°C for Exuma and 2.0°C for TOTO, $\pm 0.7^\circ\text{C}$ [Rosenheim *et al.*, 2004]. The inferred temperature increases calculated using Böhm’s [2000] $\delta^{18}\text{O}_{\text{arag}}$ – temperature calibration are only 0.9–1.0°C (Figure 2). Assuming that $\text{Sr}/\text{Ca}_{\text{sw}}$ was constant over this century, the discrepancy in secular trends between the individual proxies in each sclerosponge is interpreted as the result of salinity change. In order to estimate the salinity




Figure 2. Time series of Sr/Ca (light orange and blue) and $\delta^{18}\text{O}$ (dark orange and blue) from sclerosponges in Exuma Sound and TOTO. Temperature scales were constructed from empirical calibrations of Rosenheim *et al.* [2004] for Sr/Ca and Böhm *et al.* [2000] for $\delta^{18}\text{O}$ (constant $\delta^{18}\text{O}_{\text{sw}}$ of 0‰ SMOW is assumed). Both proxies indicate increasing temperature between 1890 and 1990 (linear regression), however the temperature change indicated by $\delta^{18}\text{O}_{\text{arag}}$, which measures salinity as well as temperature, is only half that indicated by Sr/Ca. To accommodate this difference, contemporaneous salinity change is assumed. The resulting secular salinity increases, 0.5 ± 0.34 psu in Exuma and 0.35 ± 0.32 psu in TOTO, are similar to secular salinity increases in the SMW over the past 45 y reported by both Curry *et al.* [2003] and Joyce *et al.* [1999]; however they occur over a longer time period. The large amplitude, high-frequency variability of the TOTO data set shows the effects of higher sampling resolution as well as dense plumes of hypersaline water from the Bahamas bank.

¹Auxiliary material is available at <ftp://ftp.agu.org/apend/gl/2004GL021499>.




Figure 3. 10 y smoothed salinity time series with error envelopes from each sclerosponge, in deviations from 1890–1990 mean (psu). Decadal scale variations in salinity occur simultaneously in both Exuma (blue) and TOTO (orange). The secular salinity trend observed in the SMW by *Curry et al.* [2003] is represented approximately by the white line and follows the most recent portions of both sclerosponge records of salinity. The sclerosponge salinity trends are significantly correlated to the NAO index (red line with circles, calculated in deviations from the 1890–1990 mean annual pressure difference between the Açores and Iceland (mbar) and smoothed to the 10 yr running mean). The NAO index is related to trade wind stress fluctuations as stronger trade winds develop with an elevated Açores high pressure during positive NAO events. This type of strengthened wind stress can both evaporate surface waters, changing salinity and temperature along density surfaces, as well as drive density surfaces deeper as reported by *Joyce et al.* [1999]. Despite the significance of the correlation, NAO only explains 20–30% of the salinity variance as the connections to the Bahamas are tenuous and there is still a secular component to both temperature and salinity change.

change, we combine three linear calibration equations incorporating both measured parameters ($\delta^{18}\text{O}_{\text{arag}}$ and Sr/Ca) to solve for three related unknowns ($\delta^{18}\text{O}_{\text{sw}}$, T, and S) [*Böhm et al.*, 2000; *Ganssen and Kroon*, 2000; *Rosenheim et al.*, 2004].

$$T = 19.6 - 4.9(\delta^{18}\text{O}_{\text{arag}} - \delta^{18}\text{O}_{\text{sw}}) \quad (1)$$

$$\text{Sr}/\text{Ca} = 12.5 - 0.102T \quad (2)$$

$$\delta^{18}\text{O}_{\text{sw}} = 0.54S - 19 \quad (3)$$

Oxygen isotope values ($\delta^{18}\text{O}$) are measured in ‰ VPDB for aragonite and ‰ SMOW for water. Ratios of Sr/Ca are measured in $\text{mmol}\cdot\text{mol}^{-1}$, temperature (T) is in $^{\circ}\text{C}$, and salinity (S) is in psu. Equation (3) is constructed from surface waters (<50 m) of the return flow of the N. Atlantic Gyre between 45°N and 31°N . During transport between these latitudes, this water mass begins adopting the properties of the SMW [*Krauss*, 1986; *Ottens*, 1991]. By combining and rearranging these equations to reconstruct the salinity record preserved in Bahamas sclerosponges,

$$S = \frac{1}{0.54} [40 - (2.0\text{Sr}/\text{Ca} - \delta^{18}\text{O}_{\text{arag}})] \quad (4)$$

we find secular salinity increases of 0.35 ± 0.34 psu in Exuma Sound and 0.5 ± 0.35 psu in TOTO that must happen coevally with the secular temperature increases to explain the discrepancies between Sr/Ca and $\delta^{18}\text{O}_{\text{aragonite}}$ linear temperature trends. It is evident from (4) that our reconstruction is sensitive to the gradient $\delta^{18}\text{O}_{\text{sw}}/S$ ($0.54 \text{‰}\cdot\text{psu}^{-1}$). Despite the significant cumulative residual error from equations (1)–(3) (auxiliary material), our data clearly identify salinity variation in the subsurface waters of both Exuma Sound and TOTO during the last 100 years.

4. Climate Implications

[8] Equation (4) allows the proxy reconstruction of a continuous salinity record of the SMW between 1890 and 1990, significantly improving the time resolution and extent of instrumental records. This water mass is surface forced well away from shallow shelf areas such as the Bahamas bank where local conditions can amplify climatological changes in wind stress. When such high-frequency variation is filtered from the salinity records by a 10 y running mean, several decadal-scale deviations are evident, culminating with a nearly monotonic increase in salinity over the last 40 years (Figure 3). Relative to the error inherent in equation (4) (Figure 3), the decadal-scale trends are more significant than the secular changes in salinity and are interpreted to reflect the baseline conditions of the subsurface SMW. However regional nature of secular salinity changes is suggested by similar secular Sr/Ca decreases reported in absence of coeval $\delta^{18}\text{O}_{\text{arag}}$ decreases in Jamaican sclerosponges [*Haase-Schramm et al.*, 2003].

[9] Figure 3 illustrates that the decadal-scale components of our salinity reconstructions are significantly correlated to decadal-scale variations in the NAO index between 1890–1990 ($p < 0.0005$ in TOTO and $p < 0.01$ in Exuma) supporting a recent hypothesis that monotonic salinity increase measured in the SMW is related to NAO [*Joyce et al.*, 1999; *Marshall et al.*, 2001; *Curry et al.*, 2003]. The role of the NAO in tropical and subtropical N. Atlantic variability has been suggested by its effects on the trade winds [*Marshall et al.*, 2001] via the Açores high which can strengthen the pressure gradient over which the trade winds blow. This can increase evaporation over the SMW before it is subducted from the surface and also deepen density surfaces regionally, giving the impression that surface forcing of a subsurface water mass has changed [*Joyce et al.*, 1999]. Although the oceans' response to increased heat availability is likely complex and spatially heterogeneous, the magnitude of our estimates of secular temperature increase ($1.6\text{--}2.0^{\circ}\text{C}$) compared to published global averages suggests some aspects of both surface forcing of the SMW and deepening of the Caribbean subsurface isopycnals have contributed to these trends.

[10] Coupled with the secular temperature increases of $1.6\text{--}2.0^{\circ}\text{C}$, sclerosponge salinity records add evidence to instrumental observations that regional T-S changes can surpass global averages [*Levitus et al.*, 2000; *Barnett et al.*, 2001; *Folland et al.*, 2001; *Levitus et al.*, 2001], suggesting complex, regionalized oceanic heat storage and transfer mechanisms that may be averaged by broad spatial compilations of instrumental data. Although sclerosponges from the Bahamas are limited in spatial extent, their long records of increased salinity and temperature of low-latitude

waters support previous observations that the thermohaline structure of the N. Atlantic may be adjusting to climate perturbations [Joyce *et al.*, 1999; Curry *et al.*, 2003] and these changes have been occurring some 60 years before detailed instrumental monitoring of salinity began.

[11] **Acknowledgments.** This project was supported by National Science Foundation grants 9819147 and 0136941 (to P.K.S) and 9876565 and 0134998 (to S.R.T). We thank M. Grammer and D. McNeill for assistance with the collection of the sclerosponges. W. Johns, C. Rooth, and K. Leaman gave valuable input regarding our oceanographic interpretations. G. Ganssen graciously furnished data from the APNAP cruises. This contribution benefited markedly from the constructive criticisms of two anonymous reviewers and proofreading by G. Mackenzie.

References

- Barnett, T. P., D. W. Pierce, and R. Schnur (2001), Detection of anthropogenic climate change in the world's oceans, *Science*, *292*, 270–274.
- Benavides, L. M., and E. M. Druffel (1986), Sclerosponge growth rate as determined by ^{210}Pb and $\Delta^{14}\text{C}$ chronologies, *Coral Reefs*, *4*, 221–224.
- Blanke, B., M. Arhan, A. Lazar, and G. Prévost (2002), A Lagrangian numerical investigation of the origins and fates of the salinity maximum water in the Atlantic, *J. Geophys. Res.*, *107*(C10), 3163, doi:10.1029/2002JC001318.
- Böhm, F., M. M. Joachimski, W.-C. Dullo, A. Eisenhauer, H. Lehnert, J. Reitner, and G. Woerheide (2000), Oxygen isotope fractionation in marine aragonite of coralline sponges, *Geochim. Cosmochim. Acta*, *64*, 1695–1703.
- Broecker, W. S. (1994), Massive iceberg discharges as triggers for global climate change, *Nature*, *372*, 421–424.
- Broecker, W. S., J. P. Kennet, B. P. Flower, J. T. Teller, S. Trumbore, G. Bonani, and W. Wolfli (1989), Routing of meltwater from the Laurentide Ice Sheet during the Younger Dryas cold episode, *Nature*, *341*, 318–321.
- Cardone, V. J., J. G. Greenwood, and M. A. Cane (1990), On trends in historical marine wind data, *J. Clim.*, *3*, 113–127.
- Curry, R., B. Dickson, and I. Yashahaev (2003), A change in the freshwater balance of the Atlantic Ocean over the past four decades, *Nature*, *426*, 826–829.
- Folland, C. K., et al. (2001), Global temperature change and its uncertainties since 1861, *Geophys. Res. Lett.*, *28*, 2621–2624.
- Ganssen, G. M., and D. Kroon (2000), The isotopic signature of planktonic foraminifera from NE Atlantic surface sediments: Implications for the reconstruction of past oceanic conditions, *J. Geol. Soc. London*, *157*, 693–699.
- Haase-Schramm, A., F. Böhm, A. Eisenhauer, W. Dullo, M. M. Joachimski, B. Hansen, and J. Reitner (2003), Sr/Ca ratios and oxygen isotopes from sclerosponges: Temperature history of the Caribbean mixed layer and thermocline during the Little Ice Age, *Paleoceanography*, *18*(3), 1073, doi:10.1029/2002PA000830.
- Hickey, B. M., P. MacCready, E. Elliott, and N. B. Kachel (2000), Dense saline plumes in Exuma Sound, Bahamas, *J. Geophys. Res.*, *105*, 11,471–11,488.
- Joyce, T. M., R. S. Pickart, and R. C. Millard (1999), Long-term hydrographic changes at 52 and 66°W in the North Atlantic Subtropical Gyre & Caribbean, *Deep Sea Res., Part II*, *46*, 245–278.
- Krauss, W. (1986), The North Atlantic Current, *J. Geophys. Res.*, *91*, 5061–5074.
- Lang, J. C., W. D. Hartman, and L. S. Land (1975), Sclerosponges: Primary framework constructors on the Jamaican deep fore-reef, *J. Mar. Res.*, *33*, 223–231.
- Levitus, S., J. I. Antonov, T. P. Boyer, and C. Stephens (2000), Warming of the world ocean, *Science*, *287*, 2225–2229.
- Levitus, S., J. I. Antonov, J. Wang, T. L. Delworth, K. Dixon, and A. J. Broccoli (2001), Anthropogenic warming of the Earth's climate system, *Science*, *292*, 267–270.
- Mann, M. E., and P. D. Jones (2003), Global surface temperatures over the past two millennia, *Geophys. Res. Lett.*, *30*(15), 1820, doi:10.1029/2003GL017814.
- Marshall, J., Y. Kushnir, D. Battisti, P. Chang, A. Czaja, R. Dickson, J. Hurrell, M. McCartney, R. Saravanan, and M. Visbeck (2001), North Atlantic climate variability: Phenomena, impacts and mechanisms, *Int. J. Climatol.*, *21*, 1863–1898.
- Ottens, J. J. (1991), Planktic foraminifera as North Atlantic water mass indicators, *Oceanol. Acta*, *14*, 123–140.
- Rosenheim, B. E., P. K. Swart, S. R. Thorrold, P. Willenz, L. Berry, and C. Latkoczy (2004), High resolution Sr/Ca records in sclerosponges calibrated to temperature in situ, *Geology*, *32*, 145–148.
- Schmidt, M. W., H. J. Spero, and D. W. Lea (2004), Links between salinity variation in the Caribbean and North Atlantic thermohaline circulation, *Nature*, *428*, 160–163.
- Swart, P. K., S. Thorrold, B. Rosenheim, A. Eisenhauer, C. G. A. Harrison, M. Grammer, and C. Latkoczy (2002), Intra-annual variation in the stable oxygen and carbon and trace element composition of sclerosponges, *Paleoceanography*, *17*(3), 1045, doi:10.1029/2000PA000622.
- Willenz, P., and W. D. Hartman (1999), Growth and regeneration rates of the calcareous skeleton of the Caribbean coralline sponge *Ceratoporella nicholsoni*: A long term survey, in *Proceedings of the 5th International Sponge Symposium "Origin and Outlook,"* edited by J. N. A. Hooper, *Mem. Queens. Mus.*, *44*, 675–685.
- Zhang, D., M. J. McPhaden, and W. E. Johns (2003), Observational evidence for flow between the subtropical and tropical Atlantic: The Atlantic subtropical cells, *J. Phys. Oceanogr.*, *33*, 1783–1797.

A. Eisenhauer, Department for Marine, Environmental Sciences, GEOMAR, Wischhofstraße 13, D-24118 Kiel, Germany.

B. E. Rosenheim and P. K. Swart, Division of Marine Geology and Geophysics, Rosenstiel School of Marine and Atmospheric Sciences, University of Miami, 4600 Rickenbacker Causeway, Miami, FL 33149, USA. (brosenheim@rsmas.miami.edu)

S. R. Thorrold, Biology Department MS 35, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA.

P. Willenz, Department of Invertebrates; Section Malacology, CALMARS Group, Royal Belgian Institute of Natural Sciences (IRSNB/KBIN), Rue Vautier 29, B-1000 Brussels, Belgium.