

Methane- and ammonia-oxidizing bacteria at the chemocline of Lake Kinneret (Israel)

Pilar Junier^{1,2,*}, Ok-Sun Kim^{2,3}, Werner Eckert⁴, Peter Casper⁵, Johannes F. Imhoff⁶, Karl-Paul Witzel², Ora Hadas⁴

¹Ecole Polytechnique Federale de Lausanne, Station 6, 1015 Lausanne, Switzerland

²Max-Planck Institute for Evolutionary Biology, August-Thienemann-Str. 2, 24306 Plön, Germany

³School of Biological Sciences and Institute of Microbiology, Seoul National University, 56-1 Shillim-dong, Kwanak-gu, Seoul 151-742, ROK

⁴Israel Oceanographic and Limnological Research, The Kinneret Limnological Laboratory, PO Box 447, 14950 Migdal, Israel

⁵Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Alte Fischerhütte 2, 16775 Stechlin, Germany

⁶Leibniz-Institute of Marine Sciences at the University of Kiel, Düsternbrooker Weg 20, 24105 Kiel, Germany

⁷*Present address:* Laboratory of Microbiology, University of Neuchâtel, Rue Emile-Argand 11, 2000 Neuchâtel, Switzerland

*Email: pilar.junier@unine.ch

Aquatic Microbial Ecology 58:241–248 (2010)

Fig. S1. Graphical representation of the electropherograms of pmoA/amoA PCR products digested with HaeIII. The different depths are indicated: depths 16.2 to 17.7 and Jordan River







